

Student Number	
Mark / 47	

Chemistry

The Chemical Earth and Metals Modules Test • 2003

General Instructions

- Reading time 5 minutes
- Working time 55 minutes
- Write using black or blue pen
- · Draw diagrams using pencil
- Board-approved calculators may be used
- A data sheet and a Periodic Table are provided at the back of this paper
- Write your Student Number at the top of this page

Total Marks - 47

Part A - 12 marks

- Attempt Questions 1 12
- Allow about 10 minutes for this part

Part B - 35 marks

- Attempt Questions 13 23
- Allow about 45 minutes for this part

Part A - 12 marks Attempt Questions 1-12 Allow about 10 minutes for this part

Select the alternative A, B, C or D that best answers the question. Fill in the response oval completely.

Sample: $2 + 4 = (A) \ 2 (B) \ 6 (C) \ 8 (D) \ 9$ $A \bigcirc B \bigcirc C \bigcirc D \bigcirc$

If you think you have made a mistake, put a cross through the incorrect answer and fill in the new answer.

If you change your mind and have crossed out what you consider to be the correct answer, then indicate the correct answer by writing the word **correct** and drawing an arrow as follows.



Answer Box for Questions 1–12

1	A O	ВО	c o	D O
2	A O	ВО	c o	DO
3	A O	ВО	C O	DO
4	A O	ВО	C O	DO
5	A O	ВО	\mathbf{C} O	D O
6	A O	ВО	C O	DO
7	A O	ВО	C O	DO
8	A O	ВО	C O	DO
9	A O	ВО	C O	DO
10	A O	ВО	C O	DO
11	A O	ВО	\mathbf{C} O	D O
12	A O	ВО	C O	DO

Mark your answers for Questions 1 - 12 in the Answer Box on page 2.

- 1 Which of the following describes the atomic number of an element?
 - (A) the number of protons plus neutrons in a neutral atom of the element
 - (B) the number of neutrons in a neutral atom of the element
 - (C) the number of protons in a neutral atom of the element
 - (D) the weighted mean of the isotopic masses of the element
- Which of the following was the first alloy to be used by man?
 - (A) brass
 - (B) bronze
 - (C) solder
 - (D) steel
- Which of the following describes the periodic trend in the first ionisation energy of the elements?
 - (A) decreases across a period
 - (B) increases as elements become more metallic
 - (C) decreases down a group
 - (D) increases with an increase in atomic number
- Which of the following sets is composed exclusively of covalently bonded compounds?
 - (A) Na_2O , N_2O_5 , NO
 - (B) $\text{Li}_2\text{O}, \text{Al}_2\text{O}_3, \text{P}_4\text{O}_{10}$
 - (C) NO_2 , N_2O_5 , MgO
 - (D) NO_2 , SiO_2 , P_4O_{10}
- 5 The forces holding together atoms in solid xenon would be most similar to which of these?
 - (A) the forces between the ions in NaCl
 - (B) the forces between the atoms in Cl₂
 - (C) the forces between the atoms in graphite
 - (D) the forces between the molecules of Br₂


The table shows the physical properties of three elements designated as 'X', 'Y' and 'Z'. Note that these are not the periodic table symbols of the elements.

PHYSICAL PROPERTY	Х	Y	Z
melting point (°C)	114	660	2370
electrical conductivity	nil	excellent	poor
ductile and malleable	nil	excellent	nil
metallic lustre	nil	yes	nil
density (g cm ⁻³)	4.90	2.69	2.30

Which classification of these elements given below is correct?

	X	Y	Z
(A)	non-metal	semi-metal	metal
(B)	non-metal	metal	semi-metal
(C)	semi-metal	metal	semi-metal
(D)	semi-metal	semi-metal	metal

7 The diagram represents the periodic table. The numbers represent certain elements and are not the atomic numbers for any of the elements.



Which numbers represent an element that is a semi-metal and an element that is a gas at 25°C and normal atmosphere pressure?

	Semi-metal	Gas at 25°C and normal atmosphere pressure
(A)	2	1
(B)	2	4
(C)	3	4
(D)	3	1

- Which of the following is the best description of the structure and bonding in solid iodine at room temperature?
 - (A) a network of iodine molecules with covalent bonding between the atoms, and weak intermolecular forces
 - (B) a network of iodide ions held together by a 'sea' of delocalised electrons
 - (C) a network of iodine atoms held together by covalent bonds in three dimensions
 - (D) a network of iodide ions held together by ionic bonds in a 3-dimensional lattice
- 9 The diagram below shows a block of neighbouring elements on the periodic table.

K	Ca	Sc
Rb	Sr	Y
Cs	Ва	La

Which of the following elements would have an atomic radius greater than Sr?

- (A) La, Y
- (B) Cs, Rb
- (C) Ba, Y
- (D) Ca, Sc
- The table shows the current cost of four metals which are mined and extracted in Australia.

METAL	PRICE
aluminium	\$2300 per tonne
gold	\$570 per 30 g
lead	\$750 per tonne
iron	\$1300 per tonne


Which of the following statements correctly identifies the cost factors of producing the metal?

- (A) Aluminium ore is abundant but requires much energy to extract.
- (B) Gold ore is abundant but difficult to find and extract.
- (C) Lead ore is rare but cheap to extract.
- (D) Iron ore is scarce and expensive to extract.

Which of the choices give a correct matching of the mixture and the sphere of the Earth where the mixture can be found?

sphere	Α	В	С	D
hydrosphere	seawater and fish	beach sand and gravel	seawater and fish	seawater and fish
biosphere	bacterial spores in air	seawater and fish	bacterial spores in air	seawater and fish
lithosphere	beach sand and gravel	beach sand and gravel	dusty air	dusty air
atmosphere	dusty air	bacterial spores in air	beach sand and gravel	bacterial spores in air

- (A) A
- (B) B
- (C) C
- (D) D
- 12 Study the following symbolic structures...



Which of the following best describes the substances in i, ii, iii and iv?

Substance	Α	В	C	D
i	compound	element	element	element
ii	compound	element	element	element
iii	compound	compound	element	compound
iv	compound	compound	compound	mixture

- (A) A
- (B) B
- (C) C
- (D) D

Quest	tion 13 (6 marks)
Write	balanced formulae equations for the following reactions.
(a)	calcium + water
(b)	magnesium + dilute hydrochloric acid
(c)	sodium + oxygen
Write	a pair of half-equations for the following reactions.
(d)	aluminium + dilute nitric acid
(e)	potassium + dilute sulfuric acid
(f)	barium + dilute hydrochloric acid


Question 14 (7 marks)

(a) Using the information in the table, classify the substances (A, B, C, D) as either metallic, ionic, covalent network or covalent molecular structures. (2 marks)

	SUBSTANCE			
PROPERTY	Α	В	С	D
melting point (°C)	3350	– 114	961	801
soluble in water	no	yes	no	yes
solid conducts electricity	no	no	yes	no
molten (liquid) state conducts electricity	no	no	yes	yes
hardness	very hard	solid is soft	hard	hard
other properties	crystalline solid	aqueous solution conducts	shiny solid	-

Substance A	
Substance B	
Substance C	/ ()
Substance D	

(b) Distinguish between structures C and D by drawing a diagram showing the arrangement and type of particles in each structure. (2 marks)



Question 14 continues on page 9


(c)	Explain the difference in electrical conductivity between substances C and D. (2 marks)
(d)	State the type of particles found in substance A and the type of bonding between these particles. (1 mark)
Ques	tion 15 (3 marks)
Name	e the compounds listed below.
(a)	NaHCO ₃
(b)	SF ₆
(c)	Ba(OH) ₂
Give	the formulae for the compounds listed below.
(d)	ammonium nitrate
(e)	dinitrogen pentoxide
(f)	aluminium sulfate

Question 14 (continued)

Ques	tion 16 (3 marks)
(a)	Draw the Lewis electron dot structure for the oxide ion. (1 mark)
(b)	Give the electronic configuration for the oxide ion. (1 mark)
(c)	Write the name of another species with the same electronic configuration as the oxide ion. (1 mark)
Ques	tion 17 (2 marks)
(a)	Write a balanced formulae equation for the thermal decomposition of copper(II) carbonate. (1 mark)
(b)	Describe a chemical test that can be used to prove that copper(II) carbonate can be decomposed by heat. (1 mark)
	tion 18 (1 mark) ify an example that demonstrates that light can be absorbed during the decomposition of a compound.

Question 19 (2 marks)

The diagram shows the set-up for a laboratory apparatus in which two inert (non-reactive) platinum electrodes are placed in a beaker of water that has a small amount of acid added to it.



Describe what would be observed at the electrodes when electricity flows into the the electrodes.	
Give a reason why this happens.	
Question 20 (1 mark)	
Why is the amount of energy needed to break the chemical bond in a hydrogen molecule greater than the nergy needed to break the chemical bond in a hydrogen iodide molecule?	ıe

Question 21 (4 marks)

An analytical chemist used fractional distillation to analyse a crude oil mixture from a newly discovered oil well. A 60 g sample of the crude oil was weighed accurately and then fractionally distilled. Five fractions: A, B, C, D and E were isolated and weighed. The weights obtained in grams were: 2, 18, 9, 12, 16 for components A, B, C, D and E respectively.

(a)	Construct a table of results for the data given. (1 mark)
(b)	To be an economically productive source, component B must be at least 35% by mass of the crude oil. Assess the potential economic productivity of this sample of crude oil. Justify your answer by including working. (2 marks)
(c)	Describe another situation in which gravimetric analysis supplies useful data for scientists. (1 mark)

Question 22 (1 mark)

The recycling of aluminium cans is a great success in Australia with over 50% of cans being collected and recycled! Study the equations showing the extraction and recycling of aluminium...

$$Al_2O_3 + 3C \xrightarrow{+50 \text{ MJ/kg Al}} 4Al + 3CO_2$$

$$Al_{\text{(scrap can)}} \xrightarrow{+800 \text{ kJ/kg Al}} Al_{\text{(recycled metal)}}$$

Account for the huge difference in the energy requirement	based on the nature of the processes involved.

Question 23 (5 marks)

- (a) Identify the original data which Mendeleev used to formulate the Periodic Table.
 - (i) Physical data (1 mark)
 - (ii) Chemical data (1 mark)
- (b) Historically, Mendeleev made accurate predictions about element 32 (germanium) before it was discovered. According to the HSC Periodic Table, element 117 has not yet been synthesised; however, we also can make predictions about its properties in relation to its neighbouring elements based on periodic law principles.
 - (i) Compare the relative electronegativity of element 117 with that of ununoctium (element 118).

(1 mark)

(ii) Compare the valency of element 117 with that of astatine.

(1 mark)

(iii) Compare the relative reactivity of element 117 with that of astatine.

(1 mark)

Chemistry

DATA SHEET

Avogadro's constant, N _A	$6.022 \times 10^{23} \text{ mol}^{-1}$
Volume of 1 mole ideal gas: at 101.3 kPa (1.00 atr	
	22.41 L
at 298 K (25°C)	24.47 L
Ionisation constant for water at 298 K (25°C), K_w	1.0×10^{-14}
Specific heat capacity of water	$4.18 \times 10^3 \mathrm{J kg^{-1} K^{-1}}$

Some useful formulae

 $pH = -log_{10} [H^+]$

 $\Delta H = -m C \Delta T$

Some standard potentials

		-	
$K^+ + e^-$	₹	K(s)	-2.94 V
$Ba^{2+} + 2e^{-}$	\rightleftharpoons	Ba(s)	-2.91 V
$Ca^{2+} + 2e^{-}$	₹	Ca(s)	–2.87 V
$Na^{+} + e^{-}$	\rightleftharpoons	Na(s)	–2.71 V
$Mg^{2+} + 2e^{-}$	4-7	Mg(s)	-2.36 V
$Al^{3+} + 3e^{-}$	=	Al(s)	-1.68 V
$Mn^{2+} + 2e^-$	\rightleftharpoons	Mn(s)	-1.18 V
$H_2O + e^-$	←	$\frac{1}{2}\mathrm{H}_2(g) + \mathrm{OH}^-$	–0.83 V
$Zn^{2+} + 2e^{-}$	=	Zn(s)	-0.76 V
$Fe^{2+} + 2e^{-}$	\Rightarrow	Fe(s)	-0.44 V
$Ni^{2+} + 2e^{-}$	#	Ni(s)	-0.24 V
$\mathrm{Sn}^{2+} + 2\mathrm{e}^{-}$	=	Sn(s)	–0.14 V
$Pb^{2+} + 2e^{-}$		Pb(s)	-0.13 V
$H^+ + e^-$	=	$\frac{1}{2}$ H ₂ (g)	0.00 V
$SO_4^{2-} + 4H^+ + 2e^-$	7	$SO_2(aq) + 2H_2O$	0.16 V
$Cu^{2+} + 2e^{-}$	\rightleftharpoons	Cu(s)	0.34 V
$\frac{1}{2}$ O ₂ (g) + H ₂ O + 2e ⁻	\rightleftharpoons	2OH-	0.40 V
Cu ⁺ + e ⁻	\rightleftharpoons	Cu(s)	0.52 V
$\frac{1}{2}I_2(s) + e^-$	<−	I ⁻	0.54 V
$\frac{1}{2}I_2(aq) + e^-$	\rightleftharpoons	I ⁻	0.62 V
$Fe^{3+} + e^{-}$	\rightleftharpoons	Fe ²⁺	0.77 V
$Ag^+ + e^-$	\rightleftharpoons	Ag(s)	0.80 V
$\frac{1}{2}\mathrm{Br}_2(l) + \mathrm{e}^-$	~	Br ⁻	1.08 V
$\frac{1}{2}\mathrm{Br}_2(aq) + \mathrm{e}^-$	\rightleftharpoons	Br ⁻	1.10 V
$\frac{1}{2}O_2(g) + 2H^+ + 2e^-$	~_	H_2O	1.23 V
$\frac{1}{2}\operatorname{Cl}_2(g) + \mathrm{e}^-$	\rightleftharpoons	Cl ⁻	1.36 V
$\frac{1}{2}$ Cr ₂ O ₇ ²⁻ + 7H ⁺ + 3e ⁻	_	$Cr^{3+} + \frac{7}{2}H_2O$	1.36 V
$\frac{1}{2}\text{Cl}_2(aq) + \text{e}^-$	_	CI ⁻	1.40 V
$MnO_4^- + 8H^+ + 5e^-$	< >	$Mn^{2+} + 4H_2O$	1.51 V
$\frac{1}{2}$ F ₂ (g) + e ⁻	=	F -	2.89 V

Aylward and Findlay, SI Chemical Data (4th Edition) is the principal source of data for this examination paper. Some data may have been modified for examination purposes.

H 1.008 Hydrogen 3 Li 6.941 Lithium Lithium Lithium Lithium Li 11 Na 22.99 Sodium 19 Sodium 19 K 39.10 Potassium 7 Rb 85.47 Rubidium 85.55 Cs Cs Cs St 132.9 Caesium RFrancium RFrancium RFrancium RFRANCIUM Francium RFRANCIUM RF 4 Be 9.012 Beryllium 12 Mg 24.31 Magnessium 20 Ca 40.08 Calcium 38 Sr 87.62 Strontium 56 Ba 137.3 Barium 88 Ra 1226.0] Radium 21 Sc 44.96 Scandium 39 Y 88.91 Yttrium 57–71 22 TI 47.87 Titanium 40 Zr 91.22 Zirconium 72 Zirconium 72 Hafrium 104 Rf [261.1] 23 V 50.94 Vanadium Vanadium Vanadium Vanadium Niobium 73 Ta 180.9 Tantalum 105 Db 24 Cr 52.00 Chronium 42 Mo 95.94 Molybdenu 74 W 183.8 Tungsten 106 Sg [263.1] 25 Manganess Manganess 43 43 Tc [98.91] Technetium 75 Re 186.2 Rhenium 107 Bh PERIODIC TABLE OF THE ELEMENTS Atomic Weight 26 Fe 55.85 101.1 Ruthenium 76 Os 190.2 Osmium 108 Hs [265.1] 27 Co 58.93 Cobalt 45 Rh 102.9 Rhodium 77 Ir 192.2 Iridum 109 Mt [268] 79 Au 197.0 Gold Symbol of elemen 28 Ni 58.69 Nickel 46 Pd 106.4 Palladium Palladium Platinum 110 Uun 29 Cu 63.55 Copper 47 Ag 107.9 Silver 79 Au 197.0 Gold 30 Zn 65.39 Zinc 48 Cd 112.4 Cadmium 80 Hg 200.6 Mercury 112 Uub 5 B 10.81 Boron 1.0.81 Boron 1.0.81 All 26.98 Aluminium 31 Ga 69.72 Gallium 1.14.8 Indium 1.14.8 Indium 1.14.8 Indium 1.14.8 1.11.13 6 C C 12.01 Carbon 14 Si 28.09 Silicon 272.61 Ge 72.61 Tin 118.7 Tin 82 Pb 207.2 Lead 114.4 7 N N 14.01 Nitrogen 15 P 9 30.97 30.97 33 As 74.92 Arsenic 51 Sb 121.8 Antimony 83 Bi 209.0 Bismuth 8 0 16.00 Oxygen 16 S 32.07 Sulfur Se 78.96 Selenium 52 Te 127.6 Tellurium 116 Polonium 116 Uuh 9 F 19.00 Fluorine 17 Cl 35.45 Chlorine 35 Br 79.90 Bromline 53 1 1 126.9 Iodine 85 At At 1210.0 2 He He Helium Helium 10 Ne 20.18 Nicon 18 Nicon 18 Ar 39.95 Argon Kr Kr 83.80 Krypton Krypton Krypton 131.3 Xenon Radon Radon Radon Radon Radon Radon Radon Radon Radon

Where the atomic weight is not known, the relative atomic mass of the most common radioactive isotope is shown in brackets. The atomic weights of Np and Tc are given for the isotopes ²³⁷Np and ⁹⁹Tc.

Actinides 89 Ac [227.0]

90 Th 232.0 Thorium

92 U 238.0 Uranium

93 Np [237.0] Neptunium

94 Pu [239.1]

95 Am [241.1] Americium

96 Cm [244.1]

97 Bk [249.1] Berkelium

98 Cf [252.1] Californium

100 Fm [257.1] Fermium

101 Md [258.1]

102 No [259.1] Nobelium

103 Lr [262.1] Lawrencium

Es

[252.1] Einsteinium anthanide

57 La 138.9 Lanthanum

58 Ce 140.1

59 Pr 140,9 Praseodymium

Nd Nd 144.2 Neodymiun

61 Pm [146.9] Promethium

62 Sm 150.4 Samarium

63 Eu 152.0 Europium

64 Gd 157.3 Gadoliniur

65 Tb 158.9 Terbium

66 Dy 162.5 Dysprosium

67 Ho 164.9

68 Er 167.3 Erbium

69 Tm 168.9 Thulium

70 Yb 173.0 Ytterbium

71 Lu 175.0 Lutetium