

Marking Scheme and Answers

Chemistry

Production of Materials

Theory Test • 2004

General Instructions

- Reading time – 5 minutes
- Working time – 45 minutes
- Write using black or blue pen
- Draw diagrams using pencil
- Board-approved calculators may be used
- A Data Sheet and a Periodic Table are provided at the back of this paper and may be removed for convenience
- Write your Student Number at the top of this page

Total Marks – 28

Part A – 8 marks

- Attempt Questions 1 – 8
- Allow about 15 minutes for this part

Part B – 20 marks

- Attempt Questions 9 – 12
- Allow about 30 minutes for this part

HSCFOCUS.COM

This page is intentionally blank.



Part A – 8 marks

Attempt Questions 1–8

Allow about 15 minutes for this part

Select the alternative A, B, C or D that best answers the question. Fill in the response oval completely.

Sample: $2 + 4 =$ (A) 2 (B) 6 (C) 8 (D) 9
A B C D

If you think you have made a mistake, put a cross through the incorrect answer and fill in the new answer.

A B C D

If you change your mind and have crossed out what you consider to be the correct answer, then indicate the correct answer by writing the word *correct* and drawing an arrow as follows.

A B C D
An arrow labeled "correct" points to the B option.

Answer Box for Questions 1 - 8


1	A <input checked="" type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input type="radio"/>
2	A <input type="radio"/>	B <input type="radio"/>	C <input checked="" type="radio"/>	D <input type="radio"/>
3	A <input type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input checked="" type="radio"/>
4	A <input checked="" type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input type="radio"/>
5	A <input type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input checked="" type="radio"/>
6	A <input type="radio"/>	B <input checked="" type="radio"/>	C <input type="radio"/>	D <input type="radio"/>
7	A <input type="radio"/>	B <input checked="" type="radio"/>	C <input type="radio"/>	D <input type="radio"/>
8	A <input type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input checked="" type="radio"/>

► Mark your answers for Questions 1 – 8 in the Answer Box on page 3.

1 Which of these statements describes the flow of electrons in a galvanic cell?

- (A) Electrons flow from the anode to the cathode.
- (B) Electrons flow from the cathode to the anode.
- (C) Electrons flow through the electrolyte solutions.
- (D) Electrons flow through the salt bridge between the anode and the cathode.

2 What is the IUPAC name for the compound shown below?



- (A) 2-hydroxybutane
- (B) 2-hydroxybutanol
- (C) 2-butanol
- (D) 1-methyl-1-propanol


3 Ethanol has good solubility in octane. Which statement best explains this fact?

- (A) Ethanol and octane are non-polar molecules.
- (B) Ethanol and octane are highly volatile.
- (C) Ethanol and octane both have an even number of carbon atoms.
- (D) Ethanol's ethyl group aids its solubility in octane.





4 Which of the following is the industrial source of ethylene?

- (A) cracking of alkanes
- (B) dehydration of ethanol
- (C) recycling of polyethylene
- (D) fractional distillation of crude oil

- 5 *Saran*[™] food wrap is made of an addition polymer processed into a thin, flexible cling film. A segment of the polymer molecule has the structure of...





Which of the following is the structure of the monomer?




- 6 Assuming no heat loss, what mass of ethanol must be burned to increase the temperature of 250 g of water from 25°C to 95°C, given that the heat of combustion of ethanol is 1409 kJ mol⁻¹?

- (A) 0.86 g
(B) 2.4 g
(C) 4.8 g
(D) 0.86 kg

7 Which equation shows the production of ethanol from ethylene?



8 Boris fermented a dilute solution of glucose for one week and then analysed the contents of the fermentation vessel as shown below.
Which trend describes the changes in mass during the week of fermentation?



	MASS OF...			
	CO ₂ produced	C ₂ H ₅ OH produced	C ₆ H ₁₂ O ₆	Fermentation flask
(A)	increased	increased	decreased	increased
(B)	decreased	increased	increased	increased
(C)	increased	decreased	decreased	decreased
(D)	increased	increased	decreased	decreased

Part B – 20 marks
Attempt Questions 9 – 12
Allow about 30 minutes for this part

► *Show all relevant working in questions involving calculations.*

Question 9 (5 marks)

Charlotte performs a first-hand investigation involving a galvanic cell constructed from these materials...

copper metal, 1 mol L⁻¹ copper(II) sulfate, lead metal, 1 mol L⁻¹ lead(II) nitrate, and saturated KNO₃ (aq)

- (a) Identify a hazardous risk in this experiment. (1 mark)

Lead(II) nitrate is toxic.

- (b) Identify the anode. (1 mark)

Lead

- (c) Describe the role of the salt bridge containing saturated KNO₃ solution? (1 mark)



The salt bridge completes the cell circuit.
The salt bridge allows for ion migration between the anode and cathode compartments.
The salt bridge maintains electrical charge neutrality in the anode and cathode compartments.

- (d) Charlotte lets the cell run continuously for a week. Describe TWO changes which would have occurred in the cell after one week. (2 marks)

The lead electrode becomes smaller/loses mass.
The lead(II) nitrate solution becomes more concentrated.
The copper electrode develops a coating (deposit) of copper/gains mass.
The copper(II) sulfate solution becomes less blue/less concentrated.
The cell voltage decreases.

Question 10 (4 marks)

Draw a labelled diagram of the structure of EITHER a dry cell or a lead–acid cell and write the oxidation and reduction half reactions occurring in the cell.



Dry Cell diagram should show...

- Anode 'can' of zinc. (1 mark)
- Central cathode of carbon rod surrounded by a cathodic paste of MnO_2 and carbon. (1 mark)
- Electrolyte of NH_4Cl and ZnCl_2 at the porous separator between the zinc and the cathodic paste and mixed into the cathodic paste also. (1 mark)
- Oxidation reaction... $\text{Zn}_{(s)} \rightarrow \text{Zn}^{2+}_{(aq)} + 2e^-$ (1 mark)
- Reduction reaction... $2\text{MnO}_2_{(s)} + 2\text{NH}_4^+_{(aq)} + 2\text{H}_2\text{O}_{(l)} + 2e^- \rightarrow 2\text{NH}_3_{(aq)} + 2\text{Mn}(\text{OH})_3_{(s)}$ (1 mark)
- $2\text{MnO}_2_{(s)} + 2\text{NH}_4^+_{(aq)} + 2e^- \rightarrow \text{Mn}_2\text{O}_3_{(s)} + 2\text{NH}_3_{(g)} + \text{H}_2\text{O}_{(l)}$

Lead–Acid Cell diagram should show...

- Anode plate of lead. (1 mark)
- Cathode plate of PbO_2 (1 mark)
- Electrolyte of 35% H_2SO_4 (1 mark)
- Oxidation reaction... $\text{Pb}_{(s)} + \text{SO}_4^{2-} \rightarrow \text{PbSO}_4_{(s)} + 2e^-$ (1 mark)
- Reduction reaction... $\text{PbO}_2_{(s)} + \text{SO}_4^{2-}_{(aq)} + 4\text{H}^+ + 2e^- \rightarrow \text{PbSO}_4_{(s)} + 2\text{H}_2\text{O}_{(l)}$ (1 mark)

Question 11 (5 marks)

Assess the potential of ethanol as an alternative to octane (petrol) as a car fuel.

Sample Answer

Ethanol is a renewable resource while octane is a non-renewable resource. The production and use of ethanol is carbon dioxide neutral, while petrol adds carbon dioxide to the atmosphere. Ethanol is a high octane fuel. Unlike petrol, ethanol burns cleanly and hence does not release large amounts of pollutants such as CO and aromatic hydrocarbons such as benzopyrene. As a petrol additive, it enhances the combustion of petrol. However, its production from biomass can require almost as much energy as what is obtainable from it when completely combusted. Also, being more oxygenated than petrol, it releases less energy per mole and per gram than petrol. Therefore, to obtain an equivalent amount of mileage from ethanol, more ethanol must be burnt. This requires a bigger fuel tank. The use of greater than 20% ethanol with petrol also necessitates car engine modification. There is also the problem of environmental pollution caused by the release of large quantities of fermentation liquor, soil degradation and soil erosion if vast quantities of agricultural land are devoted to crops for ethanol production.

Overall, if the production of ethanol can be made less energy demanding, such as the use of novel strains of bacteria for a more efficient fermentation, solar powered distillation units and the use of scraps and waste as raw materials, then ethanol has a very promising potential as a car fuel.

Marking Guidelines

1 – 3 Advantages cited = 1 – 3 marks

1 – 3 Disadvantages cited = 1 – 3 marks

► *At least one disadvantage must be given.*

Judgement = 1 mark

Question 12 (5 marks)

- (a) Identify a named biopolymer and the name of the specific organism or enzyme(s) used in its production. **(2 marks)**

Biopolymer name: Biopol or poly-3-hydroxybutyrate-polyhydroxy-3-valerate or poly(β -hydroxybutanoate), cellulose, cellulose nitrate, etc. **(1 mark)**

► Can be a modified natural biopolymer, e.g. rayon

Name of specific organism or enzyme(s) used in the production of the named biopolymer. **(1 mark)**

e.g. *Alcaligenes eutrophus* or bacteria. ► *Spelling errors ignored.*

- (b) Describe ONE use of the biopolymer in (a) and describe how this use (or potential use) relates to TWO properties of the biopolymer. **(3 marks)**

Use of biopolymer. **(1 mark)**

Use of biopolymer related to two properties of the biopolymer. **(2 marks)**

e.g. Biopol is used in the manufacture of shampoo bottles.

Properties related to use: Biopol is flexible, biodegradable, waterproof.

HIGHER SCHOOL CERTIFICATE EXAMINATION
Chemistry

DATA SHEET

Avogadro constant, N_A	$6.022 \times 10^{23} \text{ mol}^{-1}$
Volume of 1 mole ideal gas: at 100 kPa and	
at 0°C (273.15 K)	22.71 L
at 25°C (298.15 K)	24.79 L
Ionisation constant for water at 25°C (298.15 K), K_w	1.0×10^{-14}
Specific heat capacity of water	$4.18 \times 10^3 \text{ J kg}^{-1} \text{ K}^{-1}$

Some useful formulae

$$\text{pH} = -\log_{10}[\text{H}^+] \qquad \Delta H = -m C \Delta T$$

Some standard potentials

$\text{K}^+ + \text{e}^-$	\rightleftharpoons	K(s)	-2.94 V
$\text{Ba}^{2+} + 2\text{e}^-$	\rightleftharpoons	Ba(s)	-2.91 V
$\text{Ca}^{2+} + 2\text{e}^-$	\rightleftharpoons	Ca(s)	-2.87 V
$\text{Na}^+ + \text{e}^-$	\rightleftharpoons	Na(s)	-2.71 V
$\text{Mg}^{2+} + 2\text{e}^-$	\rightleftharpoons	Mg(s)	-2.36 V
$\text{Al}^{3+} + 3\text{e}^-$	\rightleftharpoons	Al(s)	-1.68 V
$\text{Mn}^{2+} + 2\text{e}^-$	\rightleftharpoons	Mn(s)	-1.18 V
$\text{H}_2\text{O} + \text{e}^-$	\rightleftharpoons	$\frac{1}{2}\text{H}_2(\text{g}) + \text{OH}^-$	-0.83 V
$\text{Zn}^{2+} + 2\text{e}^-$	\rightleftharpoons	Zn(s)	-0.76 V
$\text{Fe}^{2+} + 2\text{e}^-$	\rightleftharpoons	Fe(s)	-0.44 V
$\text{Ni}^{2+} + 2\text{e}^-$	\rightleftharpoons	Ni(s)	-0.24 V
$\text{Sn}^{2+} + 2\text{e}^-$	\rightleftharpoons	Sn(s)	-0.14 V
$\text{Pb}^{2+} + 2\text{e}^-$	\rightleftharpoons	Pb(s)	-0.13 V
$\text{H}^+ + \text{e}^-$	\rightleftharpoons	$\frac{1}{2}\text{H}_2(\text{g})$	0.00 V
$\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	$\text{SO}_2(\text{aq}) + 2\text{H}_2\text{O}$	0.16 V
$\text{Cu}^{2+} + 2\text{e}^-$	\rightleftharpoons	Cu(s)	0.34 V
$\frac{1}{2}\text{O}_2(\text{g}) + \text{H}_2\text{O} + 2\text{e}^-$	\rightleftharpoons	2OH^-	0.40 V
$\text{Cu}^+ + \text{e}^-$	\rightleftharpoons	Cu(s)	0.52 V
$\frac{1}{2}\text{I}_2(\text{s}) + \text{e}^-$	\rightleftharpoons	I^-	0.54 V
$\frac{1}{2}\text{I}_2(\text{aq}) + \text{e}^-$	\rightleftharpoons	I^-	0.62 V
$\text{Fe}^{3+} + \text{e}^-$	\rightleftharpoons	Fe^{2+}	0.77 V
$\text{Ag}^+ + \text{e}^-$	\rightleftharpoons	Ag(s)	0.80 V
$\frac{1}{2}\text{Br}_2(\text{l}) + \text{e}^-$	\rightleftharpoons	Br^-	1.08 V
$\frac{1}{2}\text{Br}_2(\text{aq}) + \text{e}^-$	\rightleftharpoons	Br^-	1.10 V
$\frac{1}{2}\text{O}_2(\text{g}) + 2\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	H_2O	1.23 V
$\frac{1}{2}\text{Cl}_2(\text{g}) + \text{e}^-$	\rightleftharpoons	Cl^-	1.36 V
$\frac{1}{2}\text{Cr}_2\text{O}_7^{2-} + 7\text{H}^+ + 3\text{e}^-$	\rightleftharpoons	$\text{Cr}^{3+} + \frac{7}{2}\text{H}_2\text{O}$	1.36 V
$\frac{1}{2}\text{Cl}_2(\text{aq}) + \text{e}^-$	\rightleftharpoons	Cl^-	1.40 V
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^-$	\rightleftharpoons	$\text{Mn}^{2+} + 4\text{H}_2\text{O}$	1.51 V
$\frac{1}{2}\text{F}_2(\text{g}) + \text{e}^-$	\rightleftharpoons	F^-	2.89 V

Aylward and Findlay, *SI Chemical Data* (5th Edition) is the principal source of data for this examination paper. Some data may have been modified for examination purposes.

PERIODIC TABLE OF THE ELEMENTS

		KEY									
Atomic Number	Symbol of element	Atomic Weight	Name of element	Atomic Number	Symbol of element	Atomic Weight	Name of element	Atomic Number	Symbol of element	Atomic Weight	Name of element
1	H	1.008	Hydrogen	79	Au	197.0	Gold	5	B	10.81	Boron
3	Li	6.941	Lithium	80	Hg	200.6	Mercury	6	C	12.01	Carbon
4	Be	9.012	Beryllium	81	Tl	204.4	Thallium	7	N	14.01	Nitrogen
11	Na	22.99	Sodium	82	Pb	207.2	Lead	8	O	16.00	Oxygen
12	Mg	24.31	Magnesium	83	Bi	209.0	Bismuth	9	F	19.00	Fluorine
20	Ca	40.08	Calcium	84	Po	[210.0]	Polonium	10	Ne	20.18	Neon
21	Sc	44.96	Scandium	85	At	[210.0]	Astatine	18	Ar	39.95	Argon
22	Ti	47.87	Titanium	86	Rn	[222.0]	Radon				
23	V	50.94	Vanadium	87	Fr	[223.0]	Francium				
24	Cr	52.00	Chromium	88	Ra	[226.0]	Radium				
25	Mn	54.94	Manganese	89-103	Lanthanides						
26	Fe	55.85	Iron	104	Rf	[261.1]	Rutherfordium				
27	Co	58.93	Cobalt	105	Db	[262.1]	Dubnium				
28	Ni	58.69	Nickel	106	Sg	[263.1]	Seaborgium				
29	Cu	63.55	Copper	107	Bh	[264.1]	Bhaborium				
30	Zn	65.39	Zinc	108	Hs	[265.1]	Hassium				
31	Ga	69.72	Gallium	109	Mt	[268]	Moscovium				
32	Ge	72.61	Germanium	110	Uun	—	Ununnilium				
33	As	74.92	Arsenic	111	Uuu	—	Unununium				
34	Se	78.96	Selenium	112	Uub	—	Ununbium				
35	Br	79.90	Bromine	113							
36	Kr	83.80	Krypton	114	Uuq	—	Ununquadium				
37	Rb	85.47	Rubidium	115							
38	Sr	87.62	Strontium	116	Uuh	—	Ununhexium				
39	Y	88.91	Yttrium	117							
40	Zr	91.22	Zirconium	118	Uuo	—	Ununoctium				
41	Nb	92.91	Niobium								
42	Mo	95.94	Molybdenum								
43	Tc	[98.91]	Technetium								
44	Ru	101.1	Ruthenium								
45	Rh	102.9	Rhodium								
46	Pd	106.4	Palladium								
47	Ag	107.9	Silver								
48	Cd	112.4	Cadmium								
49	In	114.8	Indium								
50	Sn	118.7	Tin								
51	Sb	121.8	Antimony								
52	Te	127.6	Tellurium								
53	I	126.9	Iodine								
54	Xe	131.3	Xenon								
55	Cs	132.9	Cesium								
56	Ba	137.3	Barium								
57	La	138.9	Lanthanum								
58	Ce	140.1	Cerium								
59	Pr	140.9	Praseodymium								
60	Nd	144.2	Neodymium								
61	Pm	[146.9]	Promethium								
62	Sm	150.4	Samarium								
63	Eu	152.0	Europium								
64	Gd	157.3	Gadolinium								
65	Tb	158.9	Terbium								
66	Dy	162.5	Dysprosium								
67	Ho	164.9	Holmium								
68	Er	167.3	Erbium								
69	Tm	168.9	Thulium								
70	Yb	173.0	Ytterbium								
71	Lu	175.0	Lutetium								
72	Hf	178.5	Hafnium								
73	Ta	180.9	Tantalum								
74	W	183.8	Tungsten								
75	Re	186.2	Rhenium								
76	Os	190.2	Osmium								
77	Ir	192.2	Iridium								
78	Pt	195.1	Platinum								
79	Au	197.0	Gold								
80	Hg	200.6	Mercury								
81	Tl	204.4	Thallium								
82	Pb	207.2	Lead								
83	Bi	209.0	Bismuth								
84	Po	[210.0]	Polonium								
85	At	[210.0]	Astatine								
86	Rn	[222.0]	Radon								
87	Fr	[223.0]	Francium								
88	Ra	[226.0]	Radium								
89-103	Lanthanides										
104	Rf	[261.1]	Rutherfordium								
105	Db	[262.1]	Dubnium								
106	Sg	[263.1]	Seaborgium								
107	Bh	[264.1]	Bhaborium								
108	Hs	[265.1]	Hassium								
109	Mt	[268]	Moscovium								
110	Uun	—	Ununnilium								
111	Uuu	—	Unununium								
112	Uub	—	Ununbium								
113											
114	Uuq	—	Ununquadium								
115											
116	Uuh	—	Ununhexium								
117											
118	Uuo	—	Ununoctium								

Lanthanides

57	La	138.9	Lanthanum
58	Ce	140.1	Cerium
59	Pr	140.9	Praseodymium
60	Nd	144.2	Neodymium
61	Pm	[146.9]	Promethium
62	Sm	150.4	Samarium
63	Eu	152.0	Europium
64	Gd	157.3	Gadolinium
65	Tb	158.9	Terbium
66	Dy	162.5	Dysprosium
67	Ho	164.9	Holmium
68	Er	167.3	Erbium
69	Tm	168.9	Thulium
70	Yb	173.0	Ytterbium
71	Lu	175.0	Lutetium

Actinides

89	Ac	[227.0]	Actinium
90	Th	232.0	Thorium
91	Pa	231.0	Protactinium
92	U	238.0	Uranium
93	Np	[237.0]	Neptunium
94	Pu	[239.1]	Plutonium
95	Am	[241.1]	Americium
96	Cm	[244.1]	Curium
97	Bk	[249.1]	Berkelium
98	Cf	[252.1]	Californium
99	Es	[252.1]	Einsteinium
100	Fm	[257.1]	Fermium
101	Md	[258.1]	Mendelevium
102	No	[259.1]	Nobelium
103	Lr	[262.1]	Lawrencium

Where the atomic weight is not known, the relative atomic mass of the most common radioactive isotope is shown in brackets.
The atomic weights of Np and Tc are given for the isotopes ²³⁷Np and ⁹⁹Tc.