



**SYDNEY BOYS HIGH SCHOOL**  
MOORE PARK, SURRY HILLS

**2004**

TRIAL HIGHER SCHOOL  
CERTIFICATE EXAMINATION


# Mathematics Extension 1

# Sample Solutions

Section	Marker
A	Mr Dunn
B	Ms Nesbitt
C	Mr Bigelow

Section A

1(a)  $(x^2 - 1)(x + 5) > 0$



AND  
 $-5 < x < -1$  (2 marks)


b)  $y = \ln \sqrt{x+1}$   
 $= \frac{1}{2} \ln(x+1)$

$$y' = \frac{1}{2(x+1)} \quad (2 \text{ marks})$$

c)  $\int_0^{\pi/6} \sec 2x \tan 2x \, dx$   
 $= \frac{1}{2} \sec 2x \Big|_0^{\pi/6}$   
 $= \frac{1}{2} \sec \frac{\pi}{3} - \frac{1}{2} \sec 0$   
 $= \frac{1}{2} \times 2 - \frac{1}{2} \times 1$   
 $= \frac{1}{2} \quad (2 \text{ marks})$

d)  $\int_0^{\sqrt{3}} \frac{dx}{9+x^2} = \frac{1}{3} \tan^{-1} \frac{x}{3} \Big|_0^{\sqrt{3}}$   
 $= \left[ \frac{1}{3} \tan^{-1} \frac{\sqrt{3}}{3} \right]$   
 $= \frac{1}{3} \tan^{-1} \frac{1}{\sqrt{3}}$   
 $= \frac{1}{3} \times \frac{\pi}{6}$   
 $= \frac{\pi}{18} \quad (2 \text{ marks})$

e) Total number of arrangements =  $7!$



If A and B are together

Then  $2 \times 6!$   
Hence not together

$$\begin{aligned} &= 7! - 2 \times 6! \\ &= 6! (7-2) \\ &= 5 \times 6! \\ &= 3600 \quad (2 \text{ marks}) \end{aligned}$$

f) LHS =  $\frac{1-\cos \theta}{\sin \theta} + \frac{\sin \theta}{1+\cos \theta}$   
 $= \frac{1-\cos^2 \theta + \sin^2 \theta}{\sin \theta (1+\cos \theta)}$

$$\begin{aligned} &= \frac{\sin^2 \theta}{\sin \theta (1+\cos \theta)} \\ &= \frac{2 \sin \theta}{1+\cos \theta} \\ \text{Let } t &= \tan \frac{\theta}{2} \\ &= \frac{2 \times \frac{2t}{1+t^2}}{1 + \frac{1-t^2}{1+t^2}} \\ &= \frac{4t}{1+t^2} = 2t \end{aligned}$$

$$= 2 \tan \frac{\theta}{2} = \text{RHS} \quad (2 \text{ marks})$$

QUESTION TWO

a)  $y = \arctan^{-1} 2x$

Let  $u = 2x$

Then  $y = \arctan^{-1} u$

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

$$= \frac{1}{\sqrt{1-u^2}} \times 2$$

$$= \frac{2}{\sqrt{1-4x^2}} \quad (2 \text{ marks})$$

b)  $y = 3 \arcsin^{-1} \sqrt{1-x^2}$

Consider  $\sqrt{1-x^2}$

$-1 \leq x \leq 1$  Range  
domain

Then

$$y = 3 \arcsin^{-1} 0 \text{ to } 3 \arcsin^{-1} 1$$

$$\therefore 0 \leq y \leq \frac{3\pi}{2} \quad \text{range} \quad (2 \text{ marks})$$

c)  $\sqrt{3} \cos x - \sin x = R \cos(x+d)$

$= R \cos x \cos d - R \sin x \sin d$

$$R \cos d = \sqrt{3}$$

$$R \sin d = 1$$

$$\tan d = \frac{1}{\sqrt{3}}$$

$$d = \frac{\pi}{6}$$

$$R^2(\cos^2 d + \sin^2 d) = 3+1$$

$$R = 2$$

c) continued

$$2 \cos\left(x + \frac{\pi}{6}\right) = 1 \quad (2 \text{ marks})$$

$$\cos\left(x + \frac{\pi}{6}\right) = \frac{1}{2}$$


$$x + \frac{\pi}{6} = \pm \frac{\pi}{3}$$

$$x = 2k\pi + \frac{\pi}{3} - \frac{\pi}{6}$$

$$x = 2k\pi + \frac{\pi}{6} \quad (2 \text{ marks})$$


$$\text{or } 2k\pi - \frac{\pi}{2}$$

d)



$\hat{F}AE = \hat{F}BC$  (angle in alternate segment)

$\hat{B}CF = \hat{C}FB$  (alternate exterior)



Hence  $\triangle ADE \sim \triangle FBA$  (2 marks)

$$\frac{EF}{EA} = \frac{EA}{EP}$$

$$EF^2 = EA \times EP$$

(3 marks)

QUESTION THREE

a) Prove  $2^{3n} - 1$  is divisible by 7 for  $n > 1$  (integer)

$$\text{Let } n=1 \text{ Then } 2^3 - 1 = 7$$

is true for  $n=1$

Assume

$$2^{3k} - 1 = 7K \text{ where } K \text{ is an integer}$$

Try to prove

$$2^{3k+3} - 1 = 7N \text{ where } N \text{ is an integer}$$

$$\text{LHS} = 2^3 \cdot 2^{3k} - 1$$

$$= 8(7K+1) - 1 \text{ from assumption.}$$

$$= 56K + 7$$

$$= 7(8K+1)$$

$$= 7N$$

True for  $n=1$

$$n=1+1=2$$

$$n=2+1=3$$

$$\vdots$$

All integers  $n > 1$  (3 marks)

$$(i) y = 1 + 2 \cos x - 2 \cos^2 x$$

$$y' = -2 \sin x + 4 \cos x \sin x$$


$$= 2 \sin x (2 \cos x - 1) \text{ (1 mark)}$$

$$(ii) y=0 \text{ when } \sin x = 0$$

$$\cos x = 1$$

$$\text{ie } x=0, \frac{\pi}{3}$$

When  $x=0, y=1$   
 $x=\frac{\pi}{3}, y=\frac{3}{2}$  } 2 marks




Max of 1.5 at  $x = \frac{\pi}{3}$

Minimum of 1 at

$$x=0 \text{ or } x=\frac{\pi}{2}$$


1 mark



$$\text{Area} = \text{bb}$$

$$= 2a \times \frac{1}{1+a^2} = \frac{2a}{1+a^2} \quad \text{(mark)}$$

$$\text{Consider } y = \frac{2x}{1+x^2}$$



$$y' = \frac{(x^2+1)2 - 2 \cdot 2x}{(1+x^2)^2}$$

$$= \frac{2-2x^2}{(1+x^2)^2}$$

$$y' = 0 \text{ when } x = \pm 1$$

THREE

c) i) continued.

When  $x = 1 + \epsilon$   $y' < 0$   
 $x = 1 - \epsilon$   $y' > 0$



Hence  $x = 1$  produces maximum

$$A_{\text{rec}} = \frac{2}{1+x} = 2 \text{ square units} \quad (\text{3 marks})$$

or

$$y'' = \frac{(1+x^4)(-4x) - (2-2x^2)4x(1+x^2)}{(1+x^2)^4}$$

$$= \frac{-4x(1+x^2)[1+x^2 + (2-2x^2)]}{(1+x^2)^4}$$

$$= \frac{-4x(1+x^2)(3-x^2)}{(1+x^2)^4}$$

When  $x = 1$   $y'' = \frac{-4 \times 2 \times 2}{2^4}$

$y'' < 0$  Hence maximum.

### QUESTION 4

$$x = -2t, \quad t = -\frac{x}{2}$$

$$y = \frac{1}{4}x^2$$

$$y' = \frac{x}{2} = -t$$

$$\text{eqn of tangent } y - t^2 = -t(x + 2t)$$

$$y - t^2 + tx + 2t^2 = 0$$

$$tx + y + t^2 = 0$$

$$tx + y + t^2 = 0$$

$$\text{at } A, y = 0$$

$$tx + t^2 = 0$$

$$t(x+t) = 0, \quad x = -t$$

$$A. (-t, 0) \quad T(-2t, t^2)$$

$$\text{Midpoint } M \left( \frac{-t-2t}{2}, \frac{0+t^2}{2} \right)$$

$$M = \left( \frac{-3t}{2}, \frac{t^2}{2} \right)$$

$$x = -\frac{3t}{2}, \quad t = -\frac{2x}{3}$$

$$y = \frac{t^2}{2}$$

$$= \frac{1}{2} \left( -\frac{2x}{3} \right)^2$$

$$y = \frac{2x^2}{9}$$

$$\text{locus of } M \quad x^2 = \frac{9}{2}y$$

$$4x^3 - 12x^2 + 11x - 3 = 0$$

roots  $\alpha - d, \alpha, \alpha + d$  (arith series)

$$\text{Sum of roots} = 3\alpha = -\frac{b}{a} = 3$$

$$\alpha = 3$$


$$\text{product } 1(1-d) + 1(1+d) + (1-d)(1+d) = \frac{c}{a}$$

$$3 - d^2 = \frac{11}{4}$$

$$d^2 = \frac{1}{4}$$

$$d = \pm \frac{1}{2}$$

$$\text{roots } \frac{1}{2}, 1, 1\frac{1}{2}.$$



$$4 \cos^2 x = \frac{1}{2}$$

$$\cos x = \frac{1}{2} \text{ or } \cos x = -\frac{1}{2}$$

$$x = -\frac{\pi}{3}, \frac{\pi}{3} \text{ or no soln in domain}$$

$$V = \pi \int_{-\frac{\pi}{3}}^{\frac{\pi}{3}} (4 \cos^2 x - \frac{1}{4} \sec^2 x) dx$$

$$2 \cos^2 x = \cos 2x + 1$$

$$V = 2\pi \int_0^{\frac{\pi}{3}} (2 \cos 2x + 2 - \frac{1}{4} \sec^2 x) dx$$

$$= 2\pi \left[ \sin 2x + 2x - \frac{1}{4} \tan x \right]_0^{\frac{\pi}{3}}$$

$$= 2\pi \left( \frac{\sqrt{3}}{2} + 2 \cdot \frac{\pi}{3} - \frac{\sqrt{3}}{4} \right) = 0$$

$$V = \left( \frac{4\pi^2}{3} + \frac{\sqrt{3}}{2} \right) u^3$$

$$(5) \text{ (a) Find } \frac{dV}{dt} = \frac{dv}{dr} \cdot \frac{dr}{dt}$$

$$\frac{dr}{dt} = -5 \text{ cm/s} \quad V = \frac{4}{3}\pi r^3$$

$$\frac{dv}{dr} = 4\pi r^2$$

$$r = 10 \text{ cm}$$

$$\frac{dv}{dt} = -5 \times 4\pi \times 10^3$$

$$= -2000\pi \text{ cm}^3/\text{s}$$

$$(b) x = 2 \cos(t + \frac{\pi}{6})$$

$$\dot{x} = -2 \sin(t + \frac{\pi}{6})$$

$$\ddot{x} = -2 \cos(t + \frac{\pi}{6})$$

$$\ddot{x} = -1^2 x, \text{ in the form } -n^2 x, n=1$$

$\therefore$  motion is SHM

$$(ii) \text{ Period} = \frac{2\pi}{n} = 2\pi$$

$$(iii) x = 2 \cos(t + \frac{\pi}{6}) = 0$$

$$t + \frac{\pi}{6} = \frac{\pi}{2} + 2n\pi$$

$$t = \frac{\pi}{3} \text{ sec (1st osc.)}$$

$$(iv) 2 \cos(t + \frac{\pi}{6}) = 1$$

$$t + \frac{\pi}{6} = +\frac{\pi}{3} + 2n\pi$$

$$t = \frac{\pi}{6} \text{ (1st osc.)}$$


$$\dot{x} = -2 \sin \frac{\pi}{3}$$

$$V = -2 \times \frac{\sqrt{3}}{2}$$

$$V = -\sqrt{3} \text{ cm/s}$$

QUESTION 5 (c)

$$\begin{aligned}
 & \int \sqrt{16-x^2} dx \\
 &= \int \sqrt{16-16\sin^2\theta} \cdot 4\cos\theta d\theta & x = 4\sin\theta \\
 & \quad \frac{dx}{d\theta} = 4\cos\theta \\
 & \quad dx = 4\cos\theta d\theta \\
 & \int \sqrt{16\cos^2\theta} \cdot 4\cos\theta d\theta \\
 & \int 4\cos\theta \cdot 4\cos\theta d\theta \\
 & 16 \int \cos^2\theta d\theta \\
 & 8 \int (\cos 2\theta + 1) d\theta & \cos 2\theta = 2\cos^2\theta - 1 \\
 & 8(\frac{1}{2}\sin 2\theta + \theta) & 2\cos^2\theta = \cos 2\theta + 1 \\
 & 4\sin 2\theta + 8\theta + C \\
 & 4 \cdot 2\sin\theta \cos\theta + 8\theta \\
 & 4 \times 2 \cdot \frac{x}{4} \frac{\sqrt{16-x^2}}{4} + 8\sin^{-1}\frac{x}{4} & \theta = \sin^{-1}\frac{x}{4} \\
 & = \frac{x}{2} \sqrt{16-x^2} + 8\sin^{-1}\frac{x}{4} + C
 \end{aligned}$$



QUESTION 6.

(a) If  $y' = \frac{3x}{4+x^2}$

$$\int y' dx = \frac{3}{2} \ln(4+x^2) + C. \quad \checkmark$$

(b).  $P(x) = 8x^3 - 12x^2 + 6x + 13$

$$P'(x) = 24x^2 - 24x + 6 \\ = 6(2x-1)^2$$

(i)  $P(x)$  is increasing where  $P'(x) > 0$ .

$$\text{i.e } 6(2x-1)^2 > 0$$

$\therefore \text{all Reals, except } x = \frac{1}{2}. \quad \checkmark$

(ii) Since  $P(x) \rightarrow -\infty$  as  $x \rightarrow -\infty$ ,  $P(0) = 13$ .  
and  $P(x)$  is increasing for all  $x \neq \frac{1}{2}$ .  
it follows that there must be a root  $x_1$  where  $x_1 < 0$ .

(iii)  $a_2 = a_1 - \frac{f(a_1)}{f'(a_1)}$

$$\text{if } a_1 = -1 \text{ then } a_2 = -1 - \frac{-8-12-6+13}{24+24+6} \\ = -1 - \frac{-13}{54} \\ = -1 + \frac{13}{54} \\ = \boxed{-0.76} \text{ (2.D.P). } \quad \checkmark$$

$$(i) T = S + A e^{-kt} \quad \text{--- (A)}$$

$$\begin{aligned}\therefore \frac{dT}{dt} &= -kA e^{-kt} \\ &= -k(T-S) \text{ from (A)}\end{aligned}$$

(ii) When  $t=0$ ,  $T=1390$  and  $S=30$  (constant)

$$\therefore 1390 = 30 + A e^0$$

$$\therefore A = 1360.$$

$$\therefore T = 30 + 1360 e^{-kt}$$

When  $t=10$ ,  $T = 1060$ .

$$\therefore 1060 = 30 + 1360 e^{-10k}$$

$$\frac{1030}{1360} = e^{-10k}$$

$$-10k = \ln \frac{103}{136}$$

$$k \doteq 0.0278.$$

$$\therefore T = 30 + 1360 e^{-0.0278t}$$

$$\text{Let } T = 110. \quad -0.0278t$$

$$110 = 30 + 1360 e^{-0.0278t}$$

$$\frac{80}{1360} = e^{-0.0278t}$$


$$\ln \frac{8}{136} = -0.0278t$$

$$t = \frac{\ln \frac{1}{17}}{-0.0278}$$

$$\doteq 102 \text{ mins}$$

$\therefore$  [it takes 92 mins longer.]

(11)



QUESTION 7.

(a) now

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots + \binom{n}{r}x^r + \dots + \binom{n}{n}x^n \quad (A)$$

(i) Differentiate both sides of (A) above -

$$n(1+x)^{n-1} = \binom{n}{1} + 2\binom{n}{2}x + 3\binom{n}{3}x^2 + \dots + r\binom{n}{r}x^{r-1} + \dots + n\binom{n}{n}x^{n-1}$$

Let  $x = 1$

$$n \cdot 2^{n-1} = \binom{n}{1} + 2\binom{n}{2} + 3\binom{n}{3} + \dots + r\binom{n}{r} + \dots + n\binom{n}{n}$$

$$n \cdot \left[ \sum_{r=1}^n r\binom{n}{r} \right] = n \cdot 2^{n-1} \quad \text{NB This is equivalent to } \sum_{r=0}^n r\binom{n}{r} = n \cdot 2^{n-1}.$$

(ii) R.T.P.  $\sum_{r=0}^n (r+1)\binom{n}{r} = 2^{n-1}(n+2)$

$$\begin{aligned} LHS &= \sum_{r=0}^n r\binom{n}{r} + \sum_{r=0}^n \binom{n}{r} \\ &= n \cdot 2^{n-1} + 2^n \quad \left( \begin{array}{l} \text{if we let } x=1 \\ \text{in (A)} \end{array} \right) \\ &= \boxed{2^{n-1}(n+2)} \quad \text{VV} \quad 2^n = \sum_{r=0}^n \binom{n}{r} \\ &= RHS. \end{aligned}$$

$$(b) (i) x = vt \Rightarrow t = \frac{x}{v}$$

$\therefore y = -\frac{1}{2}gt^2 + h$ . becomes

$$y = -\frac{1}{2}g\left(\frac{x}{v}\right)^2 + h.$$

$$\boxed{y = h - \frac{1}{2}g \frac{x^2}{v^2}} \quad \checkmark$$

$$(ii) x = vt \cos \alpha \Rightarrow t = \frac{x}{v \cos \alpha} \quad \therefore y = -\frac{1}{2}gt^2 + vt \sin \alpha + h.$$

$$\text{becomes } y = -\frac{1}{2}g\left(\frac{x}{v \cos \alpha}\right)^2 + v \frac{x \sin \alpha}{v \cos \alpha} + h.$$

$$\text{ie } \boxed{y = x \tan \alpha - \frac{gx^2 \sec^2 \alpha}{2v^2} + h} \quad \checkmark$$

(iii) Substitute  $(d, 0)$  in (i)  $0 = h - \frac{gd^2}{2v^2}$

$$\therefore h = \boxed{\frac{gd^2}{2v^2}} \quad \checkmark$$

(iv) Substitute  $(d, 0)$  in (ii).

$$0 = d \tan \alpha - \frac{gd^2}{2v^2} \sec^2 \alpha + h.$$

$$0 = d \tan \alpha - h(1 + \tan^2 \alpha) + h \quad (h = \frac{gd^2}{2v^2})$$

$$\therefore h \tan^2 \alpha - d \tan \alpha = 0$$

$$\tan \alpha (h \tan \alpha - d) = 0$$

$$\therefore \tan \alpha = 0 \text{ or } \tan \alpha = \frac{d}{h}$$

$$\text{Clearly } \tan \alpha \neq 0 \quad \therefore \boxed{\tan \alpha = \frac{d}{h}} \quad \checkmark$$

(v) Substitute  $(2d, 0)$  into (ii).

$$2d \tan \alpha - \frac{g \cdot 4d^2 \sec^2 \alpha}{2\sqrt{r}} + h = 0.$$

$$2d \tan \alpha - 4h \sec^2 \alpha + h = 0.$$

$$2d \tan \alpha - 4h(1 + \tan^2 \alpha) + h = 0$$

$$2d \tan \alpha - 4h - 4h \tan^2 \alpha + h = 0$$

$$4h \tan^2 \alpha - 2d \tan \alpha + 3h = 0$$

For  $\tan \alpha$  to be real  $4d^2 - 4 \times 4h \times 3h \geq 0$ .

$$\text{i.e. } 4d^2 - 48h^2 \geq 0.$$

$$4d^2 \geq 48h^2$$

$$d^2 \geq 12h^2$$

$$| d \geq 2h\sqrt{3} | \quad \checkmark$$