

QUESTION 1.

(a)  $y = -x$

$m_1 = -1$

$y = \frac{1}{\sqrt{3}}x$

$m_2 = \frac{1}{\sqrt{3}}$

ie/  $m_2 = \frac{\sqrt{3}}{3}$

$$\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$$

$$= \left| \frac{-1 - \frac{\sqrt{3}}{3}}{1 - \frac{\sqrt{3}}{3}} \right|$$


$$= \frac{3 + \sqrt{3}}{3 - \sqrt{3}}$$

$\therefore \theta = 75^\circ$


(b)  $(2x+1)(x-1) > 3(x-1)^2$

$(x-1)[2x+1-3(x-1)] > 0$

$(x-1)(4-x) > 0$



$\therefore 1 < x < 4$



$l = r\theta$

$= 8 \times \frac{3\pi}{4}$

$= 6\pi$

$\therefore$  the circumf. is  $6\pi$  cm.

(d) (i)  $f(-x) = \frac{8}{4+(-x)^2}$

$= \frac{8}{4+x^2}$

$= f(x)$


$\therefore f(x)$  is an even fn.

(ii)  $\lim_{x \rightarrow \infty} \frac{8}{4+x^2} = \lim_{x \rightarrow \infty} \frac{8/x^2}{4/x^2+1}$

$= \frac{0}{0+1}$

$= 0$

(iii) when  $x=0$ ,  $y=2$



QUESTION 2.

a)  $x = 76$  (alt. seg. thm)

b) reflex  $\angle AOC = 234^\circ$  (angle at centre is twice  $\angle$  at circumf.)

$$\theta = 234^\circ - 180^\circ \text{ (COB is a diam)}$$

$$\therefore \theta = 54^\circ$$

c)  $\angle WXY = 90^\circ$  ( $\angle$  in a semi-circle)

$$\angle OXY = 37^\circ \text{ (base } \angle \text{'s of isos. } \Delta \text{)}$$

$$\therefore \angle WXD = 53^\circ$$

d) i) In  $\Delta PRT$  and  $\Delta QPT$

$LT$  is common

$$\angle TPR = \angle TPQ \text{ (alt. seg. thm)}$$

$\therefore \Delta PRT \sim \Delta QPT$  (equiangular)

ii)  $\frac{PT}{QT} = \frac{RT}{PT}$  (ratio of corres. sides in sim.  $\Delta$ 's)

$$\therefore PT^2 = QT \times RT$$

e) ii)  $BW = BX$  (tangents to a circle, from ext. pt., are equal)

$$AW = AZ \quad " \quad " \quad "$$

$$DY = DZ \quad " \quad " \quad "$$

$$CY = CX \quad " \quad " \quad "$$

So,  $(AW + BW) + (DY + CY) = (AZ + BX) + (DZ + CX)$

ie  $AB + DC = (AZ + DZ) + (BX + CX)$

$$= AD + BC$$

$$\therefore AB + DC = AD + BC$$

### QUESTION 3

(a) Let  $P(x) = 2x^3 + Kx^2 - 18x - 8$ , then  $P(-2) = 0$

$$0 = -16 + 4K + 36 - 8$$


$$\therefore K = -3$$

$$\begin{array}{r} 2x^2 - 7x - 4 \\ x+2 \overline{) 2x^3 - 3x^2 - 18x - 8} \\ \underline{2x^3 + 4x^2} \phantom{- 8} \\ -7x^2 - 18x \phantom{- 8} \\ \underline{-7x^2 - 14x} \phantom{- 8} \\ -4x - 8 \\ \underline{-4x - 8} \\ 0 \end{array}$$

$$\therefore 2x^3 - 3x^2 - 18x - 8 = (x+2)(x-4)(2x+1)$$

(b)  $P(x) = x(x+1)^2(2x-1)$

$$P(1) > 0$$



$$\begin{aligned} \text{(c) (i) } \alpha + \beta + \gamma &= -b/a \\ &= -2 \end{aligned}$$

$$\begin{aligned} \text{(ii) } \alpha\beta + \alpha\gamma + \beta\gamma &= c/a \\ &= 3 \end{aligned}$$

$$\begin{aligned} \text{(iii) } \alpha^2 + \beta^2 + \gamma^2 &= (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \alpha\gamma + \beta\gamma) \\ &= 4 - 6 \\ &= -2 \end{aligned}$$

(d) Let the roots be  $\alpha, \alpha, \beta$

$$\text{then } 2\alpha + \beta = 6 \quad \text{--- (1)}$$

$$\alpha^2 + 2\beta\alpha = 9 \quad \text{--- (2)}$$

$$\alpha^2\beta = K \quad \text{--- (3)}$$

$$\text{from } \textcircled{1} \quad \beta = 6 - 2\alpha$$

$$\text{sub. in } \textcircled{2} \quad \alpha^2 + 2\alpha(6 - 2\alpha) = 9$$

$$\alpha^2 + 12\alpha - 4\alpha^2 = 9$$

$$-3\alpha^2 + 12\alpha - 9 = 0$$

$$\alpha^2 - 4\alpha + 3 = 0$$

$$(\alpha - 3)(\alpha - 1) = 0$$

$$\alpha = 3, 1$$

$$\text{when } \alpha = 3, \quad \beta = 0 \quad \text{and } k = 0$$

$$\text{" } \alpha = 1, \quad \beta = 4 \quad \text{and } k = 4$$

$\therefore$  if  $k = 0$ , the roots are 3, 3, 0


if  $k = 4$ , " " " " 1, 1, 4

#### QUESTION 4

$$\begin{aligned} \text{e) } \tan A + \cot B &= \frac{\sin A}{\cos A} + \frac{\cos B}{\sin B} \\ &= \frac{\sin A \sin B + \cos A \cos B}{\cos A \sin B} \\ &= \frac{\cos(A - B)}{\cos A \sin B} \end{aligned}$$

$$\begin{aligned} \cot A + \tan B &= \frac{\cos A}{\sin A} + \frac{\sin B}{\cos B} \\ &= \frac{\cos A \cos B + \sin A \sin B}{\sin A \cos B} \\ &= \frac{\cos(A - B)}{\sin A \cos B} \end{aligned}$$

$$\begin{aligned} \text{LHS} &= \frac{\cos A \sin B}{\cos(A - B)} + \frac{\sin A \cos B}{\cos(A - B)} \\ &= \frac{\cos A \sin B + \sin A \cos B}{\cos(A - B)} \\ &= \frac{\sin(A + B)}{\cos(A - B)} \\ &= \text{RHS.} \end{aligned}$$



(b) Let  $y = \cos^4 x$

$$\frac{dy}{dx} = 4 \cos^3 x \times -\sin x$$

ie/  $\frac{dy}{dx} = -4 \sin x \cos^3 x$

(c)  $7 \sin \theta + \cos \theta = R \sin(\theta + \alpha)$

$$R = \sqrt{50}$$

ie  $R = 5\sqrt{2}$

$$\tan \alpha = 1/7$$


$$\alpha = 8^\circ 8'$$

$$5\sqrt{2} \sin(\theta + 8^\circ 8') = 5$$

$$0^\circ < \theta \leq 360^\circ$$

$$\sin(\theta + 8^\circ 8') = 1/\sqrt{2}$$

$$8^\circ 8' \leq \theta \leq 368^\circ 8'$$



$$\theta + 8^\circ 8' = 45^\circ \text{ or } 135^\circ$$

$$\theta = 36^\circ 52' \text{ or } 126^\circ 52'$$

or //  $7 \cdot \frac{2t}{1+t^2} + \frac{1-t^2}{1+t^2} = 5$   $0^\circ \leq \frac{\theta}{2} \leq 180^\circ$

$$14t + 1 - t^2 = 5 + 5t^2$$

$$6t^2 - 14t + 4 = 0$$


$$3t^2 - 7t + 2 = 0$$

$$(3t-1)(t-2) = 0$$

$$\tan \theta/2 = 1/3 \text{ or } 2$$

$$\theta/2 = 18^\circ 26' \text{ or } 63^\circ 26'$$

$$\therefore \theta = 36^\circ 52' \text{ or } 126^\circ 52'$$



$$\begin{aligned}
 (d) \int_0^{\pi/9} \sec 3x \tan 3x \, dx &= \frac{1}{3} \left[ \sec 3x \right]_0^{\pi/9} \\
 &= \frac{1}{3} \left[ \sec \pi/3 - \sec 0 \right] \\
 &= \frac{1}{3} (2 - 1) \\
 &= \frac{1}{3}
 \end{aligned}$$

### QUESTION 5

$$\begin{aligned}
 (a) \text{ Eqn of chord PQ: } m &= \frac{ap^2 - aq^2}{2ap - 2aq} \\
 &= \frac{a(p-q)(p+q)}{2a(p-q)} \\
 &= \frac{p+q}{2}
 \end{aligned}$$

$$y - ap^2 = \frac{p+q}{2} (x - 2ap)$$

Since PQ is a focal chord, it passes through the focus  $(0, a)$

$$a - ap^2 = \frac{p+q}{2} (0 - 2ap)$$

$$a - ap^2 = -ap(p+q)$$

$$a - ap^2 = -ap^2 - apq$$

$$a = -apq$$

$$\therefore pq = -1$$

$$(b) \quad i) \quad y = x^2 / 4a$$

$$\frac{dy}{dx} = x / 2a$$

$$\text{at } T, \quad m = 2at / 2a \\ = t$$

$$y - at^2 = t(x - 2at)$$

$$y - at^2 = tx - 2at^2$$

$\therefore y - tx + at^2 = 0$ , is the eq<sup>n</sup> of the tangent at T.

ii) when  $y = 0$ ,  $x = at \quad \therefore A$  is the point  $(at, 0)$

when  $x = 0$ ,  $y = -at^2 \quad \therefore B$  is the point  $(0, -at^2)$

iii) Show that  $\Delta STB$  is isosceles

$$SB = a + at^2 \\ = a(1 + t^2)$$

$$ST^2 = (2at - 0)^2 + (at^2 - a)^2 \\ = 4a^2t^2 + a^2t^4 - 2a^2t^2 + a^2 \\ = a^2t^4 + 2a^2t^2 + a^2 \\ = a^2(t^4 + 2t^2 + 1) \\ = a^2(t^2 + 1)^2$$

$$ST = a(t^2 + 1)$$

$$\therefore SB = ST$$

$\therefore$  the tangent is equally inclined to the y-axis and the focal chord through T.

$$(iv) \quad A(x_1, y_1) \quad B(x_2, y_2) \quad k : l \quad T(x, y)$$

$$2at = \frac{k \cdot 0 + l \cdot at}{k + l}$$

$$at^2 = \frac{-kat^2 + l \cdot 0}{k + l}$$


\* or, use  
the diag.

$$k + l = \frac{lat}{2at} \\ = \frac{l}{2}$$

$$k + l = \frac{-kat^2}{at^2} \\ = -k$$

$$\therefore \frac{l}{2} = -k \\ \text{ie/ } \frac{k}{l} = -\frac{1}{2}$$

$\therefore T$  divides  $AB$  externally  
in the ratio  $1 : 2$



QUESTION 6.

$$a) \log_3 \left( \frac{9x-2}{x^2} \right) = 2$$

$$\therefore \frac{9x-2}{x^2} = 3^2$$

$$9x-2 = 9x^2$$

$$9x^2 - 9x + 2 = 0$$

$$(3x-2)(3x-1) = 0$$

$$\therefore x = \frac{2}{3}, \frac{1}{3}$$

$$(b) \int_0^1 \frac{3x^2}{1+x^3} dx = \left[ \log(1+x^3) \right]_0^1$$

$$= \log 2 - \log 1$$

$$= \log 2$$

$$(c) V = 2\pi \int_0^1 (e^{2x} + 2 + e^{-2x}) dx$$

$$= 2\pi \left[ \frac{e^{2x}}{2} + 2x - \frac{e^{-2x}}{2} \right]_0^1$$

$$= 2\pi \left[ \frac{e^2}{2} + 2 - \frac{e^{-2}}{2} - \left( \frac{1}{2} - \frac{1}{2} \right) \right]$$

$$y = e^x + e^{-x}$$
$$y^2 = e^{2x} + 2 + e^{-2x}$$

$\therefore$  vol. is  $4\pi (e^2 + 4 - \frac{1}{e^2})$  cub. units.

$$(d) y = xe^{-x}$$

$$u = x$$

$$v = e^{-x}$$

$$u' = 1$$

$$v' = -e^{-x}$$

$$\frac{dy}{dx} = e^{-x} - xe^{-x}$$

$$\frac{d^2y}{dx^2} = -e^{-x} - (e^{-x} - xe^{-x})$$

$$= -e^{-x} - e^{-x} + xe^{-x}$$

$$= e^{-x} (x-2)$$

St. pts. occur when  $y' = 0$

$$\text{i.e. } e^{-x} (1-x) = 0$$

$$\therefore x = 1$$



when  $x=1$ ,  $y = e^{-1}$ ,  $y'' < 0$

$\therefore$  there is a max. turning point at  $(1, 1/e)$

ii) Inflexion occurs when  $y'' = 0$  and concavity changes.

$$y'' = 0 \quad \text{when } x=2, \quad y = 2e^{-2}$$


when  $x=1$ ,  $y'' < 0$

"  $x=3$ ,  $y'' > 0$

$\therefore$  concavity has changed.

$(2, 2/e^2)$  is a point of inflexion

iii)



QUESTION 7

$$\begin{aligned} \text{a) } \int_0^4 x(16-x^2)^{\frac{1}{2}} dx &= -\frac{1}{2} \int_0^4 -2x(16-x^2)^{\frac{1}{2}} dx \\ &= -\frac{1}{2} \cdot \frac{2}{3} \left[ (16-x^2)^{\frac{3}{2}} \right]_0^4 \\ &= -\frac{1}{3} \left[ 0 - 16^{\frac{3}{2}} \right] \\ &= 64/3 \end{aligned}$$

$$\begin{aligned} \text{OR // } \int_0^4 (16-x^2)^{\frac{1}{2}} \cdot x dx &= -\frac{1}{2} \int_{16}^0 u^{\frac{1}{2}} du \\ &= -\frac{1}{2} \cdot \frac{2}{3} \left[ u^{\frac{3}{2}} \right]_{16}^0 \\ &= -\frac{1}{3} \left[ 0 - 16^{\frac{3}{2}} \right] \\ &= 64/3 \end{aligned}$$

$$u = 16 - x^2$$

$$du = -2x dx$$

$$x=0, \quad u=16$$

$$x=4, \quad u=0$$

$$\begin{aligned}
 \text{b) } V &= \pi \int_0^{\pi/3} \tan^2 x \, dx \\
 &= \pi \int_0^{\pi/3} (\sec^2 x - 1) \, dx \\
 &= \pi \left[ \tan x - x \right]_0^{\pi/3}
 \end{aligned}$$

$$\sin^2 x + \cos^2 x = 1$$

$$\tan^2 x + 1 = \sec^2 x$$

$$\begin{aligned}
 &= \pi \left[ \tan \frac{\pi}{3} - \frac{\pi}{3} - (\tan 0 - 0) \right] \\
 &= \pi \left( \sqrt{3} - \frac{\pi}{3} \right)
 \end{aligned}$$

$\therefore$  volume is  $\frac{\pi}{3}(3\sqrt{3} - \pi)$  cub. units.

$$\text{c) } \cos 2x = \cos^2 x - \sin^2 x$$

$$\cos 2x = 1 - 2\sin^2 x$$

$$2\sin^2 x = 1 - \cos 2x$$

$$\text{ie/ } \sin^2 x = \frac{1}{2}(1 - \cos 2x)$$

$$\begin{aligned}
 \int_0^{\pi/2} \sin^2 \frac{\theta}{2} \, d\theta &= \frac{1}{2} \int_0^{\pi/2} (1 - \cos \theta) \, d\theta \\
 &= \frac{1}{2} \left[ \theta - \sin \theta \right]_0^{\pi/2} \\
 &= \frac{1}{2} \left[ \frac{\pi}{2} - \sin \frac{\pi}{2} - (0 - \sin 0) \right] \\
 &= \frac{1}{2} \left[ \frac{\pi}{2} - 1 \right] \\
 &= \frac{1}{4} (\pi - 2)
 \end{aligned}$$

$$\text{d) } f''(x) = -18 \cos 3x$$

$$f'(x) = -6 \sin 3x + c$$

$$f'(x) = 0 \text{ when } x = \frac{2\pi}{3}$$

$$0 = -6 \sin 2\pi + c$$

$$\text{ie/ } c = 0$$

$$f(x) = 2 \cos 3x + k$$

$$1 = 2 \cos 2\pi + k$$

$$\text{ie/ } k = -1$$

$$\therefore f(x) = 2 \cos 3x - 1$$