



TRINITY GRAMMAR SCHOOL  
MATHEMATICS DEPARTMENT



YEAR 12 2008 HALF YEARLY EXAMINATIONS

# MATHEMATICS EXTENSION 1

## (3 UNIT COMPONENT)

ASSESSMENT TASK 3

WEIGHTING 30%

Examination Date:

Friday 2<sup>nd</sup> May 2008

OUTCOMES REFERRED TO: HE1, HE2, HE3, HE4, HE6, HE7, PE1, PE2, PE3, PE4, PE6.

### General Instructions

- Reading time – **5 minutes**.
- Working time – **2 hours**.
- Write using blue or black pen.
- Board-approved calculators may be used.
- A table of standard integrals is provided at the back of this paper.
- All necessary working should be shown in every question.
- **Begin** each question in a **new booklet**.
- Write your **examination number** and your **teacher's name** on the front of each answer booklet.

### Total marks – 84

- Attempt Questions 1-7.
- All questions are of equal value.
- Mark values are shown at the side of each question part.

HSCFOCUS.COM

**BLANK PAGE**

**QUESTION 1** (12 marks) Use a SEPARATE writing booklet.

**Marks**

- (a) Solve the inequation  $\frac{4}{5-x} \leq 1$ . 3
- (b) Use the substitution  $u = 1 + \tan x$  to evaluate  $\int_0^{\frac{\pi}{4}} \frac{\sec^2 x}{\sqrt{1 + \tan x}} dx$ . 3
- (c) Find the following:
- i)  $\int \frac{3x}{x^2 + 9} dx$ . 2
- ii)  $\int \frac{3}{x^2 + 9} dx$ . 2
- iii)  $\int \frac{x^2 + 9}{3x} dx$ . 2

**QUESTION 2** (12 marks) Use a SEPARATE writing booklet.

**Marks**


- (a) Solve  $\frac{x+1}{x^2+1} > 1$ . 2
- (b) Find  $\int x(1-x^2)^5 dx$ , using the substitution  $u = 1 - x^2$ , or otherwise. 2
- (c) Find the acute angle, correct to the nearest minute, between the lines  $3x + y = 4$  and  $x - y = 1$ . 2
- (d) The point  $P(19, -15)$  divides an interval  $AB$  externally in the ratio  $3 : 2$ . Find the coordinates of the point  $B(x, y)$  given  $A(-2, 3)$ . 3
- (e) Prove by mathematical induction that if  $n$  is a positive integer, then: 3
- $$\frac{1}{1 \times 5} + \frac{1}{5 \times 9} + \frac{1}{9 \times 13} + \dots + \frac{1}{(4n-3)(4n+1)} = \frac{n}{4n+1}$$

**QUESTION 3** (12 marks) Use a SEPARATE writing booklet.

**Marks**

- (a) i) If  $f(x) = e^{x+2}$ , find the inverse function  $f^{-1}(x)$ . 2
- ii) On the same axes, sketch the graphs of  $y = f(x)$  and  $y = f^{-1}(x)$ . 2
- (b) Differentiate  $x^2 \cos^{-1}(3x)$ . 2
- (c) Find the exact value of  $\cos^{-1}\left(\sin \frac{4\pi}{3}\right)$ . 3
- (d) Determine the domain and range of  $y = 2 \sin^{-1}\left(\frac{x}{3}\right)$  and hence sketch the graph. 3

(a)



In the diagram, the points  $A, B$  and  $O$  are in the same horizontal plane.

$A$  and  $B$  are 50m apart and  $\angle AOB = 60^\circ$ .  $OT$  is a vertical tower of height  $h$  metres.

The angles of elevation of  $T$  from  $A$  and  $B$  respectively are  $45^\circ$  and  $\alpha$ .

( $\alpha$  is acute.)

- i) Show  $AO = h$ . 1
  - ii) Show  $OB = h \cot \alpha$ . 1
  - iii) By using the cosine rule in triangle  $AOB$ , show that: 1  

$$h^2 \cot^2 \alpha - h^2 \cot \alpha + h^2 = 50^2.$$
  - iv) Given the tower is 30 high, find the angle  $\alpha$  correct to the nearest degree. 2
- (b) Write down the general solution, in terms of  $\pi$ , of the equation  $\tan \theta = -\frac{1}{\sqrt{3}}$ . 2
- (c)  $\alpha$  and  $\beta$  are acute angles such that  $\cos \alpha = \frac{3}{5}$  and  $\sin \beta = \frac{1}{\sqrt{5}}$ . 2  
 Without finding the size of either angle, show that  $\alpha = 2\beta$ .
- (d) Without using a calculator, find the exact value of  $\sin\left(\cos^{-1}\frac{2}{3} + \sin^{-1}\frac{1}{4}\right)$ . 3

**QUESTION 5** (12 marks) Use a SEPARATE writing booklet.

**Marks**

- (a) Newton's Law of Cooling states that when an object at temperature  $T$  ( $^{\circ}\text{C}$ ) is placed in an environment at a temperature  $R$  ( $^{\circ}\text{C}$ ), then the rate of temperature loss is given by the equation  $\frac{dT}{dt} = k(T - R)$ ; where  $t$  is the time in seconds and  $k$  is a constant.

A packet of peas, initially at  $24^{\circ}\text{C}$  is placed in a snap-freeze refrigerator in which the internal temperature is maintained at  $-40^{\circ}\text{C}$ . After 5 seconds the temperature of the packet is  $19^{\circ}\text{C}$ . Suppose  $T = R + Ae^{kt}$ , where  $A$  is a constant.

- i) State the value of  $A$ . 2
- ii) Show that  $k = \frac{1}{5} \log_e \left( \frac{59}{64} \right)$ . 2
- iii) Hence show that it will take approximately 29 seconds for the packet's temperature to reduce to  $0^{\circ}\text{C}$ . 2

**Question 5 continues on page 7.**

**QUESTION 5 (continued)**

(b) If  $y = \frac{1}{200} t e^{-t}$ , show that  $\frac{dy}{dt} = \frac{1}{200} (1-t) e^{-t}$ . **1**

- (c) Kiran has recently consumed three standard alcoholic drinks. Immediately after he has finished his last drink, his blood alcohol level is measured over a four-hour period.

Let his blood alcohol level at any time  $t$  be  $A$ , where  $t$  is the time in hours after his last drink.


It is found that the rate of change  $\frac{dA}{dt}$  of his blood alcohol content is given by:

$$\frac{dA}{dt} = \frac{1}{200} (1-t) e^{-t}, \text{ where } 0 \leq t \leq 4.$$

- i) Show that his blood alcohol content increases during the first hour and decreases after the first hour. **2**
- ii) Initially his blood alcohol content was  $0.0005$ . Find  $A$  as a function of  $t$ . **2**  
You will need to use part (b).
- iii) Determine his maximum alcohol content during the four-hour period. **1**  
Give your answer correct to four decimal places.

**End of Question 5.**


(a)



In the diagram above, two circles touch one another externally at the point  $A$ .  
 A straight line through  $A$  meets one of the circles at  $T$  and the other at  $S$ .  
 The tangents at  $T$  and  $S$  meet the common tangent at  $A$  at  $X$  and  $Y$  respectively.  
 Let  $\theta = \angle XTA$ .

- i) Explain why  $\angle XAT$  is  $\theta$ . 1
- ii) Prove that  $TX \parallel YS$ . 2

(b)



$ABCD$  is a cyclic quadrilateral in which  $AB = AC$ , and  $CD$  is produced to  $E$ .  
 Prove that  $AD$  bisects  $\angle BDE$ . (Hint: let  $\angle ADE = \theta$ ).


3

Question 6 continues on page 9.



**QUESTION 6 (continued)**

(c)



In the diagram above,  $ABP$  is a triangle inscribed in a circle.

The altitudes  $BN$  and  $AM$  of the triangle intersect at  $H$ .

The altitude  $AM$  is produced to meet the circumference of the circle at  $F$ .

Copy the diagram into your examination booklet.

Let  $\angle PBF = \alpha$ .

- |      |   |   |
|------|---|---|
| i)   | Why is $\angle PAF = \alpha$ ?                    | 1 |
| ii)  | Why are points $A$ , $N$ , $M$ and $B$ concyclic? | 1 |
| iii) | Why is $\angle NBM = \alpha$ ?                    | 1 |
| iv)  | Show that $M$ bisects $HF$ .                      | 3 |

**End of Question 6.**

**QUESTION 7** (12 marks) Use a SEPARATE writing booklet.

**Marks**

(a) Two points  $P(2ap, ap^2)$  and  $Q(2aq, aq^2)$  lie on the parabola  $x^2 = 4ay$  ( $a > 0$ ).

i) By derivation, show that the equation of the chord is: **2**

$$y = \frac{1}{2}(p+q)x - apq.$$

ii) If the chord  $PQ$  passes through the focus,  $S$ , show that  $pq = -1$ . **1**


iii) Using the fact that  $PQ = PS + SQ$ , or otherwise, show that **3**

the chord  $PQ$  has length  $a\left(p + \frac{1}{p}\right)^2$ .

**Question 7 continues on page 11.**

**QUESTION 7 (continued)**

(b)



The tangent at  $T(2t, t^2)$ ,  $t \neq 0$ , on the parabola  $x^2 = 4y$  meets the  $x$ -axis at  $A$ .

$P(x, y)$  is the foot of the perpendicular from  $A$  to  $OT$ , where  $O$  is the origin.

The equation of the tangent at  $T$  is  $y = tx - t^2$ .

- i) Prove that the equation of  $AP$  is  $y = -\frac{2}{t}(x - t)$ . 2
- ii) Show that the equation of  $OT$  is  $t = \frac{2y}{x}$ . 1
- iii) Hence, or otherwise, prove that the locus of  $P(x, y)$  lies on a circle with centre  $(0, 1)$  and give its radius. 3

**End of paper.**

## STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left( x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left( x + \sqrt{x^2 + a^2} \right)$$

NOTE :  $\ln x = \log_e x, \quad x > 0$