



**SYDNEY BOYS HIGH
SCHOOL**
MOORE PARK, SURRY HILLS

2009

**TRIAL HIGHER SCHOOL
CERTIFICATE EXAMINATION**

Mathematics Extension 1

General Instructions

- Reading Time – 5 Minutes
- Working time – 120 Minutes
- Write using black or blue pen. Pencil may be used for diagrams.
- Board approved calculators may be used.
- Start each **NEW** question in a separate answer booklet.
- Marks may **NOT** be awarded for messy or badly arranged work.
- All answers must be given in exact simplified form unless otherwise stated.
- All necessary working should be shown in every question.

Total Marks – 84

- Attempt questions 1-7.

Examiner: *D.McQuillan*

This is an assessment task only and does not necessarily reflect the content or format of the Higher School Certificate.

START EACH NEW QUESTION ON A NEW ANSWER BOOKLET

Question 1 (12 marks)

Marks

(a) Solve $x(3 - 2x) > 0$.

2

(b) Find $\frac{d}{dx}(e^{-x} \cos^{-1} x)$

2

(c) The remainder when $x^3 + ax^2 - 3x + 5$ is divided by $(x + 2)$ is 11. Find the value of a .

2

(d) Find the general solution of $2 \cos x + \sqrt{3} = 0$.

2

(e) Solve $\frac{x^2 - 9}{x} \geq 0$.

2

(f) Find $\int_0^2 (4 + x^2)^{-1} dx$.

2

End of Question 1

START EACH NEW QUESTION ON A NEW ANSWER BOOKLET

Question 2 (12 marks)

Marks

- (a) Use the substitution $x = \ln u$ to find $\int \frac{e^x}{\sqrt{1-e^{2x}}} dx$. **3**
- (b) Use one application of Newton's method to find an approximation to the root of the equation $\cos x = x$ near $x = 0.5$. Give your answer correct to two decimal places. **3**
- (c) The curves $y = e^{2x}$ and $y = 1 + 4x - x^2$ intersect at the point $(0, 1)$. Find the angle between the two curves at this point of intersection. **3**
- (d) **3**
- (i) In how many ways can a committee of 2 Englishmen, 2 Frenchmen and 1 American be chosen from 6 Englishmen, 7 Frenchmen and 3 Americans.
- (ii) In how many of these ways do a particular Englishman and a particular Frenchman belong to the committee?

End of Question 2

START EACH NEW QUESTION ON A NEW ANSWER BOOKLET

Question 3 (12 marks)

Marks

- (a) Evaluate $\cos\left(\sin^{-1}\left(-\frac{1}{2}\right)\right)$. 1
- (b) 3
- (i) Expand $\cos(\alpha + \beta)$.
- (ii) Show that $\cos 2\alpha = 1 - 2\sin^2 \alpha$.
- (iii) Evaluate $\lim_{x \rightarrow 0} \frac{1 - \cos 2x}{x^2}$.
- (c) If $\alpha = \tan^{-1}\left(\frac{5}{12}\right)$ and $\beta = \cos^{-1}\left(\frac{4}{5}\right)$, calculate the exact value of $\tan(\alpha - \beta)$. 2
- (d) A and B are points $(-1, 7)$ and $(5, -2)$; P divides AB internally in the ratio $k : 1$. 3
- (i) Write down the coordinates of P in terms of k .
- (ii) If P lies on the line $5x - 4y = 1$, find the ratio of AP:PB.
- (e) Use mathematical induction to prove that 3
- $$1 \times 1! + 2 \times 2! + 3 \times 3! + \cdots + n \times n! = (n+1)! - 1,$$
- where n is a positive integer.

End of Question 3


START EACH NEW QUESTION ON A NEW ANSWER BOOKLET

Question 4 (12 marks)

Marks

- (a) If $\frac{dy}{dx} = 1 + y$ and when $x = 0$, $y = 2$ find y as a function of x . **3**

- (b) Two circles cut at A and B. A line through A meets one circle at P and the other at Q. BR is a tangent to circle ABP and R lies on circle ABQ. Prove that $PB \parallel QR$. **3**



- (c) The area bounded by the curve $y = \sin^{-1} x$ the y axis and $y = \frac{\pi}{2}$ is rotated about the y axis. Find the volume of the solid generated. **3**

- (d) A particle moves in a straight line from a position of rest at a fixed origin O. Its velocity is v when displacement from O is x . If its acceleration is $\frac{1}{(x+3)^2}$, find v in terms of x . **3**

End of Question 4

START EACH NEW QUESTION ON A NEW ANSWER BOOKLET

Question 5 (12 marks)

Marks

- (a) The speed $v \text{ ms}^{-1}$ of a particle moving along the x axis is given by $v^2 = 24 - 6x - 3x^2$, where $x \text{ m}$ is the distance of the particle from the origin. **4**
- (i) Show that the particle is executing Simple Harmonic Motion.
- (ii) Find the amplitude and the period of motion.
- (b) Five Jovians and four Martians are sitting around discussing galactic peace. **5**
- (i) In how many ways can they be arranged around the table?
- (ii) If Marvin the Martian will not sit next to any of the Jovians, how many arrangements are possible?
- (iii) If all the Jovians sit together and all the Martians sit together and Marvin will still not sit next to a Jovian, how many arrangements are possible?
- (c) If one root of $x^3 + px^2 + qx + r = 0$ equals the sum of the two other roots, prove that $p^3 + 8r = 4pq$. **3**


End of Question 5

START EACH NEW QUESTION ON A NEW ANSWER BOOKLET

Question 6 (12 marks)

Marks

- (a) $f(x) = \cos x - \sqrt{3} \sin x$, where $0 \leq x \leq 2\pi$. **3**
- (i) Write $f(x)$ in the form $R \cos(x + \alpha)$ where $R > 0$ and α is in the first quadrant.
- (ii) Hence solve $f(x) = 1$.
- (b) Wheat falls from an auger onto a conical pile at the rate of $20 \text{ cm}^3 \text{ s}^{-1}$. The radius of the base of the pile is always equal to half its height. **5**
- (i) Show that $V = \frac{1}{12} \pi h^3$ and hence find $\frac{dh}{dt}$.
- (ii) Find the rate at which the pile is rising when it is 8 cm deep, in terms of π .
- (iii) Find the time taken for the pile to reach a height of 8 cm.
- (c) In a horizontal triangle APB, $AP = 2AB$, and the angle A is a right angle. On AB stands a vertical and regular hexagon ABCDEF. Prove that PC is inclined to the horizontal at an angle whose tangent is $\frac{\sqrt{3}}{5}$. **4**



End of Question 6

START EACH NEW QUESTION ON A NEW ANSWER BOOKLET

Question 7 (12 marks)

Marks

- (a) Use mathematical induction to prove that $\cos(\pi n) = (-1)^n$, where n is a positive integer. 2

- (b) 3
- (i) Find the largest possible domain of positive values for which $f(x) = x^2 - 5x + 13$ has an inverse.

- (ii) Find the equation of the inverse function, $f^{-1}(x)$.

- (c) The straight line $y = mx + b$ meets the parabola $x^2 = 4ay$ at the points $P(2ap, ap^2)$ and $Q(2aq, aq^2)$. 7

- (i) Find the equation of the chord PQ and hence or otherwise show that $pq = -\frac{b}{a}$.

- (ii) Prove that $p^2 + q^2 = 4m^2 + \frac{2b}{a}$.

- (iii) Given that the equation of the normal to the parabola at P is $x + py = 2ap + ap^3$ and that N, the point of intersection of the normals at P and Q, has coordinates

$$\left[-apq(p+q), a(2 + p^2 + pq + q^2) \right],$$

express these coordinates in terms of a , m and b .

- (iv) Suppose that the chord PQ is free to move while maintaining a fixed gradient. Find the locus of N and show that this locus is a straight line.

Verify that this line is a normal to the parabola.

End of Question 7

End of Exam

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax,$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE: $\ln x = \log_e x, \quad x > 0$