

JRAHS 2005 TRIAL HSC – EXTENSION I

Question 1. [Start a new page] Marks

- a) If $P(x) = x^3 - 2x^2 + ax + 4$ is divisible by $(x + 2)$, what is the value of a ? 1
- b) i) Find $\frac{d}{dx} \ln(\cos 2x)$ 1
- ii) Hence evaluate exactly $\int_0^{\frac{\pi}{6}} \tan 2x \, dx$ 2
- c) Find i) $\int \frac{e^{3x} dx}{2 + e^{3x}}$ 1
- ii) $\int \frac{dx}{\sqrt{9 - 4x^2}}$ 2
- d) Find the acute angle between the straight lines $y = \sqrt{3}x + 2$ and $x = 2$. 2
- e) Solve : $x + 2 < \frac{4}{x - 1}$ ($x \neq 1$) 3

Question 2. [Start a new page] Marks

- a) By making the substitution $u = \sqrt{x}$, evaluate exactly $\int_0^{\frac{\pi^2}{16}} \frac{\sin \sqrt{x}}{\sqrt{x}} dx$ 3
- b) i) Sketch the graph of the curve $y = 3 \sin^{-1}(x/2)$, clearly indicating the domain and range. 2
- ii) Find the area enclosed between the curve $y = 3 \sin^{-1}(x/2)$, the line $y = (3\pi/2)$ and the positive y axis. 2
- c) The polynomial equation $3x^3 - 2x^2 + 3x - 4 = 0$ has roots α, β and γ . 2
Find the exact value of $\frac{1}{\alpha\beta} + \frac{1}{\beta\gamma} + \frac{1}{\gamma\alpha}$.
- d) The letters of the word **MOUSE** are to be rearranged.
- i) How many arrangements are there which start with the letter **M** and end with the letter **E**? 1
- ii) How many arrangements are there in which the vowels are grouped together? (A vowel is one of the letters **A, E, I, O, U**) 1

- iii) How would your answers to parts (i) and (ii) change if the given word had been **MOOSE** instead of **MOUSE**? 1

Question 3. [Start a new page] **Marks**

- a) Find the general solution (in radian form) of the equation $\cos 2x = \cos x$ 3

- b) i) At the distinct points $P(2at, at^2)$ and $Q(2au, au^2)$ on the parabola $4ay = x^2$, the tangents are drawn. You may assume, without proof, that the equation of the tangent at P is $y = tx - at^2$. Show that the tangents from P and Q intersect at the point $(a(u + t), aut)$. 2

- ii) From the point R $(a, -6a)$, two tangents are drawn to the parabola $4ay = x^2$. If the points of contact of these tangents are P and Q, show that the triangle PQR is isosceles. 3

- c) Suppose that $(5 + 2x)^{12} = \sum_{k=0}^{12} a_k x^k$.

- i) Using the Binomial Theorem, write an expression for a_k . 2

- ii) Show that $\frac{a_{k+1}}{a_k} = \frac{24 - 2k}{5k + 5}$ 2

Question 4. [Start a new page] **Marks**

- a) i) Sketch the function $y = f(x)$ where $f(x) = (x - 2)^2 - 4$, clearly showing all intercepts on the axes. (Use the same scale on both axes) 2

- ii) What is the largest positive domain of f for which $f(x)$ has a continuous inverse $f^{-1}(x)$? 1

- iii) Sketch the graph of $f^{-1}(x)$ on the same axes as (i). 1

- b) A particle moves along the x axis according to the equation $x = 6 \sin 2t - 2\sqrt{3} \cos 2t$.

- i) Express x in the form $R \sin(2t - \alpha)$ where $R > 0$ and $0 \leq \alpha \leq \pi/2$. 2

- ii) Prove that the particle moves in simple harmonic motion. 1

- c) A, B and C are three sequential points on a straight line on horizontal ground. A vertical flagpole PQ is situated close by the line (but its base P is not on the line). 5

The angles of elevation of the top of the flagpole from A, B and C are $\tan^{-1} \frac{1}{4}$, $\tan^{-1} \frac{1}{2}$ and $\tan^{-1} \frac{1}{3}$ respectively. If $AB = 90\text{m}$ and $BC = 30\text{m}$, find the height of the flagpole.

Question 5. [Start a new page]

Marks


- a) A particle is moving along the x -axis. Its velocity, v m/s at position x metres is given by

$$v = \sqrt{5x - x^2}.$$

Find the acceleration of the particle when $x = 2$.

- b) Prove by induction that, for any positive integer n , the product $(n+1)(n+2)\dots(n+n)$ is always a multiple of 2^n but never a multiple of 2^{n+1} .

c)



In the diagram, CQ and BP are altitudes of the triangle ABC . The lines CQ and BP intersect at T , and AT is produced to meet CB at R .

- i) Prove that $\angle TAQ = \angle QCB$. 3
- ii) Prove that $AR \perp CB$. 2

Question 6. [Start a new page]

Marks

- a) Cane sugar, when placed in water, converts into dextrose at a rate which is proportional to the amount of unconverted material remaining. That is, if M grams is the amount of material converted after t minutes, then

$$dM/dt = k(S - M)$$

where S grams is the initial amount of cane sugar and k is a constant.


- i) Show that $M = S + Ae^{-kt}$ satisfies the equation, where A is a constant. 1
- ii) If a certain amount of cane sugar is placed in water at time $t = 0$ and 40% of it has been converted after 10 minutes, show that the value of k is $\frac{1}{10} \log_e \left(\frac{5}{3}\right)$. 2
- iii) How long will it take, to the nearest minute, for 99% of the cane sugar to be converted into dextrose? 2

(Question 6 is continued on the next page)

Question 6. (Continued)

Marks

b)



The diagram above shows two particles *A* and *B* projected from the origin. particle *A* is projected with initial velocity V m/s at an angle α and particle *B* is projected T seconds later with the same initial velocity V m/s but an angle of β . The particles collide at the point *R*.

i) You may assume that the equation of the path of *A* is given by

$$y = -\frac{gx^2}{2V^2} \sec^2 \alpha + x \tan \alpha$$

Write down the equation of the path of *B*.

1

Show that the x-coordinate of the collision point *R* is given by

$$x = \frac{2V^2 \cos \alpha \cos \beta}{g \sin(\alpha + \beta)}$$

3

ii) You may assume that the horizontal displacement of *A* after t seconds is given by

$$x = Vt \cos \alpha$$

(a) Write down the equation for the horizontal displacement of *B* (Remember that *B* is projected T seconds after *A*).

1

(β) Show that, for the collision to take place, the value of T is given by

$$T = \frac{2V(\cos \beta - \cos \alpha)}{g \sin(\alpha + \beta)} \quad 2$$

Question 7. [Start a new page]

Marks

a) It is known that 5% of all gear boxes made in Factory A are faulty whereas 7% of gear boxes made in Factory B are faulty. If 120 gear boxes are bought, 10 from each factory, what is the probability that exactly two are faulty? 4

b) i) By rotating the circle $x^2 + y^2 = r^2$ about the x axis between appropriate limits, show that the volume V of a spherical cap of height h , as shown in Figure 1, is given by

$$V = \frac{\pi h^2}{3}(3r - h) \quad (0 \leq h \leq 2r) \quad 3$$




Figure 1




Figure 2

A chemical flask is modeled by surmounting an open cylinder on a thin spherical shell (with a matching circular opening at the top). See Figure 2.

ii) The body of the flask is of radius 8cm. The neck has radius $2\sqrt{7}$ cm. and height 16cm. Show that the total height of the flask is 30cm. 1

iii) Water is poured into the flask at a constant rate of $100 \text{ cm}^3/\text{sec}$. If h is the depth of the water in the flask, use the result from part (i) to find an expression (in terms of h) for the rate at which the water level rises in the spherical portion of the flask. 2

iv) Find this rate at the instant when the water level reaches the base of 2

the cylinder and hence, or otherwise, calculate how long it will take (from that point in time) to overflow the flask. Give your answer to the nearest second.

THIS IS THE END OF THE EXAMINATION

HSCFscGuides.net.com