### JRAHS 2006 TRIAL HSC - EXT I

## **Ouestion 1**.

(a) Solve for x: 
$$\frac{1}{x-2} \ge 2$$
.

(b) Find: 
$$\lim_{h\to 0} \left( \frac{\cos 2h - 1}{h} \right)$$
.

- The point P divides A(-1, 5) and B(3, -2) in the ratio r:1. (c) Find the coordinates of *P* in terms of *r*. (i)
  - Find the value of *r* when the line 2x 3y + 4 = 0 intersects (ii) the interval *AB*.
- $\int_{0}^{1} (x^2 + 1)^3 dx.$ (d) Evaluate

#### [START A NEW PAGE] **Question 2**.

A plate is initially heated to  $55^{\circ}$  C, and it then cools to  $41^{\circ}$  C in 10 minutes. If the surrounding temperature,  $S^{\circ}$  C, is  $22^{\circ}$  C and assuming Newton's Law of Cooling: (a)

$$\frac{dT}{dt} = -k(T-S).$$

- Find the temperature of the plate 25 minutes from the 3 (i) start of cooling (to 1 decimal place).
- Find the time for the plate to cool to  $25^0 C$  (to 1 decimal place). 2 (ii)
- Sketch the graph of the rate of temperature,  $\frac{dT}{dt}$ , versus the temperature T. 1 (iii)

(b) The displacement x metres of a particle after t seconds, is given by:  $x = 5\sin 3t - 7\cos 3t.$ 

- Show that the motion of the particle is SHM. (i) 2 (ii) Find the maximum displacement. 1 Find the time when the particle first passes through the centre of 2 (iii) motion (correct to 1 decimal place).
  - Sketch the graph of the acceleration  $\ddot{x}$  versus displacement x. 1 (iv)

Marks

3

2

2

2

Question 3. [START A NEW PAGE]

(a) Differentiate 
$$\cos^{-1}\left(-\frac{1}{x}\right)$$
 with respect to x. Answer in simplified form. 3


(b) (i) On the same set of axes, sketch the graphs of  $y = \sin^{-1} x$  and  $y = \tan^{-1} x$ . 2

- (ii) Given that:  $\int_{0}^{1} \sin^{-1} x dx = \frac{\pi}{2} 1$ , find the area of the region bounded by **3**  $y = \sin^{-1} x, y = \tan^{-1} x$  and x = 1.
- (c) (i) Show that  $y = e^{-x} \sin 2x$  is a solution to the differential equation: **3** y'' + 2y' + 5y = 0.
  - (ii) Hence, or otherwise, find  $\int e^{-x} \sin 2x \, dx$ .

1

# Question 4. [START A NEW PAGE]

(a) A fire truck arrives at a burning building 10 metres high and 15 metres wide. The water nozzle hose on the fire truck is 2 metres above the ground and *d* metres from the building, as shown in the diagram.



The angle of elevation of the hose,  $\alpha$ , can be adjusted to range from  $10^0$  to  $45^0$ . The parametric equations for the water particles from the nozzle are given by:  $x = 30t \cos \alpha$  and  $y = 30t \sin \alpha - 5t^2$ , where t is the time in seconds when g = 10.

(i) Show that the trajectory path of the water is given by the equation:

$$y = x \tan \alpha - \frac{x^2}{180} (1 + \tan^2 \alpha).$$

(ii) The hose nozzle is adjusted to an angle of elevation of  $45^{\circ}$ . 2 Find the distance, *d*, from the building if the water is to reach the furthest point *B* on top of the building as shown (answer to the nearest centimetre). Q 4 continues over the page

Q 4 part (a) <i>continued</i> Mar			
	(iii)	Find the angle of elevation $\alpha$ of the nozzle, for the water to reach position <i>A</i> , when the hose nozzle is 20 metres from the burning building (answer to nearest minute).	2
(b)	Find	$\int \frac{4x-7}{2x^2+1} dx.$	3
(c)	(i)	For $t > 0$ , find the limiting sum of: $e^{-t} + e^{-2t} + e^{-3t} + \dots$	1
	(ii)	Hence, find an expression for the series; $e^{-t} + 2e^{-2t} + 3e^{-3t} + \dots$	1
(d)	A sem	ni-circle of radius <i>r</i> has the equation: $y = \sqrt{r^2 - x^2}$ .	
	(i)	Find $\frac{dy}{dx}$ at the point $P(x, y)$ .	1
	(ii)	Prove that the tangent, at any point <i>P</i> on the semi-circle, is perpendicular to the radius.	1
Question 5.		[START A NEW PAGE]	
(a)	Find t	Find the greatest coefficient in the expansion of $(4x+5)^{11}$ .	
	(Leave the answer in index form).		
(b)	A ping pong ball is initially placed 1 metre beneath the surface of the water,		
	as sho	we in the diagram. $x$	

0 water surface

The ping pong ball is released in the water with an acceleration of  $\ddot{x} \text{ m/s}^2$ , where  $\ddot{x} = -625x$ , and where x metres is the displacement of the motion measured from the water surface.

(i) Is the motion of the ping pong ball only SHM? Give reasons.	1
(ii) Prove that: $\frac{d}{dx}\left(\frac{v^2}{2}\right) = \ddot{x}.$	2
(iii) Find the expression for the ping pong ball's velocity $v$ m/s	2
when it is in the water.	
(iv) Find the velocity of the ball at the water's surface.	1
(v) Assuming there is no air resistance and the acceleration due to gravity	2
is 10 m/s <sup>2</sup> , derive an expression for the displacement in air in terms of	v
(vi) Find the maximum height that the ping pong ball reaches above the surface of the water.	1

#### Question 6. [START A NEW PAGE]

- (a) How many groups of 2 men and 2 women can be hosen from 6 men and 8 women?
- (b) Six letter words are formed from the letters of the word *CYCLIC*.
  - (i) How many different 6-letter words can be formed?
  - (ii) How many 6 letter words can be formed, if no 'C's are together?
  - (iii) What is the probability of all the 'C's together, if it is known a vowel is 2 at the end?
- (c) Prove, by the method of mathematical induction that:  $\sin q + \sin 3q + \sin 5q + ... + \sin(2n-1)q = \frac{1 - \cos 2nq}{2\sin q}$ , for n = 1, 2, 3, ...

## Question 7. [START A NEW PAGE]

- (a) At the end of each month, for 15 years, a man invests \$400 at an interest rate Which is paid monthly at 6% *pa*.
  - (i) Show that the value of his first payment, at the end of 15 years, is \$976.75
  - (ii) Find the value of the man's total investment at the end of the 15 years. 2
- (b) A circle, centre *O* with a constant radius *r*, is such that the chords *AC* and *BD* intersect at point *E*,  $\angle CED = \theta$  radians and  $\angle BOC = \frac{2\pi}{3}$  radians, as shown the diagram.

s shown the diagram.

Not to scale

Show that the sum of the arcs AB and CD equal  $2r\theta$ , give reasons. 3

0

(ii) Show that the perimeter P of the shape ABCD, where BC, AD are chords 2 and CD, AB are arc lengths, is given by:

$$P = r\left(2\theta + \sqrt{3} + 2\sin\left(\frac{\pi}{3} - \theta\right)\right)$$

C

(iii) Find the value of  $\theta$ , in the domain  $0 \le \theta \le \frac{\pi}{2}$  for the perimeter of *ABCD* **3** to have a maximum value. Justify your answer.

(i)

Marks

2

4