

2002
Higher School Certificate
Trial Examination

Chemistry

General Instructions

- Reading time 5 minutes
- Working time 2½ hours
- Board-approved calculators may be used
- Write using black or blue pen
- Draw diagrams using pencil
- A Data Sheet and Periodic Table are provided at the back of this paper
- Write your student number and/or name at the top of every page

Total marks - 85

This section has two parts, Part A and Part B

Part A

Total marks (15)

- Attempt Questions 1–15
- Allow about 30 minutes for this part

Part B


Total marks (70)

- Attempt Questions 16-31
- Allow about 2 hours for this part

This paper MUST NOT be removed from the examination room

A zinc rod is placed in four different solutions, as shown in the diagrams below.

1.



You would notice a displacement reaction in beakers

- (A) 1 and 2
- (B) 1 and 3
- (C) 1 and 4
- (D) 2 and 3
- 2. Bromine water can be used to distinguish between:
 - (A) ethane and propane
 - (B) propane and propene
 - (C) ethene and propene
 - (D) ethane and ethanol
- 3. Which of the following metals could reduce $iron(\Pi)$ ions in aqueous solution?
 - (A) zinc
 - (B) tin
 - (C) copper
 - (D) silver
- 4. Consider the reaction described by the equation below:

$$C_2H_5OH \rightarrow C_2H_4 + H_2O$$

This reaction is an example of:

- (A) polymerisation
- (B) hydration
- (C) dehydration
- (D) addition

5. The table below gives some properties of radioactive isotopes

Name	Radiation emitted	Half Life
Americium-241	α	432 years
Carbon-14	β	5.7x10 ³ years
Magnesium-27	β	9.5 minutes
Sodium-24	β,γ	15 hours

The isotope that is most suitable for detecting leaks in water pipes would be.

- (A) Americium-241
- (B) Carbon-14
- (C) Magnesium-27
- (D) Sodium-24

Questions 6 and 7 refer to the table below which shows the colour ranges of three acid-base indicators.

Indicator	Ve	Very acidic Neutral				Ver	Very basic								
pН	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Bromothymol blue	yel	low				***	> 6.0		7.6←				t	olue	
Methyl orange	red		-	→ 3.1	4.5	5←			у	ellow	1				
Phenolphthalein	col	ourle	ess						→8.3	3	10.0)←		red	

- 6. A solution is yellow in bromothymol and methyl orange, and colourless in phenolphthalein. What is the pH range of the solution?
 - (A) 7.5 to 8.5
 - (B) 6.0 to 7.5
 - (C) 4.5 to 6.0
 - (D) 8.5 to 10.0
- 7. 0.1 mol L⁻¹ citric acid (C₆H₈O₇) solution is neutralised with a solution of 0.1 mol L⁻¹ sodium hydroxide (NaOH). The best indicator for this titration would be:
 - (A) methyl orange
 - (B) phenolphthalein
 - (C) a mixture of methyl orange and bromothymol blue.
 - (D) bromothymol blue

8.

In the upper atmosphere, the most important radiation absorbed by ozone is

- (A) microwaves
- (B) infra-red
- (C) light
- (D) ultra-violet

9. The pH of four acids and their concentrations are shown in the table below.

Acid	Conc. (mol L-1)	pH
A	0.1	1.0
В	0.05	1.0
С	0.01	2.0
D	0.1	2.0

Which acid in the table is the weakest?

- (A) A
- (B) B
- (C) C
- (D) D

10. The pH of a solution of magnesium hydroxide of concentration 4.5×10^{-3} mol L⁻¹ is closest to:

- (A) 11.9
- (B) 11.6
- (C) 2.1
- (D) 2.4

11. What is the correct systematic name for the following compound?


- (A) 1,1,2,2,2 difluorotrichloroethane
- (B) ethane -1,1 difluoride -2,2,2 trichloride
- (C) 1,1,1 trichloro 2,2 difluoroethane
- (D) trichlorodifluoroethane

12. A student analysed a sample of water from an underground spring which was suspected to be contaminated with certain metallic ions.

The student added dilute hydrochloric acid to the solution and noted no changes. She then added dilute ammonia to a second sample and also noted no change. She then performed a flame test and noted a faint but distinct red colour.

The ion in the water is most probably:

- (A) Fe³⁺
- (B) Ba^{2+}
- (C) Cu²⁺
- (D) Ca^{2+}
- 13. The diagram below represents the stages in the treatment of sewage.



Secondary treatment chlorination would be used to:

- (A) adjust the pH
- (B) improve the BOD levels
- (C) kill pathogens
- (D) activate enzymes necessary for sewage digestion

STUDENT NUMBER/NA	ME:
-------------------	-----

- 14. The technique best suited to detecting low concentrations of metal ions in solutions would be:
 - (A) fractional distillation
 - (B) gas-liquid chromatography (GLC)
 - (C) atomic absorption spectroscopy (AAS)
 - (D) nuclear magnetic resonance spectroscopy (NMR)
- 15. The following table shows the potential release of acid into waterways from some minerals as a result of mining.

Potential acid release from some alunite group minerals					
Mineral	Formula	moles H ⁺ /mole			
Plumbojarosite	PbFe ₆ (SO ₄) ₄ (OH) ₁₂	6			
Benverite	Pb(Cu,Fe,Al) ₃ (SO ₄) ₂ (OH) ₆	0-5			
Natrojarosite	NaFe ₃ (SO ₄) ₂ (OH) ₆	3			
Alunite	KAl ₃ (SO ₄) ₂ (OH) ₆	0			

The most noticeable effect of run off from a mine waste heap with a high concentration of plumbojarosite into a nearby stream would be:

- (A) a rapid rise in BOD levels
- (B) pH levels would drop
- (C) dissolved oxygen levels would drop
- (D) eutrophication would begin

	STUDENT NUMBER/NAME:	*** *
Secti	ion I - continued	
Atte	t B al Marks (70) ampt Questions 1 – 32 aw about 2 hours for this part.	
Ansv	wer the questions in the spaces provided	
Show	v all relevant working in questions involving calculations.	
CMA-		Marks
Many which	y cosmetic and pharmaceutical preparations require the use of a solvent such as ethanolic h can be produced by the fermentation of sugars.	1,
(a)	Relate the use of ethanol as a solvent to the nature of the ethanol molecule.	2
(b)	Describe the conditions required to produce ethanol by the fermentation of glucose.	2
(c)	Explain how mass changes can be used to monitor the fermentation reaction. A suitable equation should be included in your answer.	2